

From axioms to synthetic inference rules via focusing

Sonia Marin

Dept. of Computer Science, University College London, UK

Dale Miller

Inria-Saclay & LIX, Ecole Polytechnique, Palaiseau, France

Elaine Pimentel

Departamento de Matemática, UFRN, Brazil

Marco Volpe

Fortiss, Munich, Germany

Abstract

An important application of *focused* variants of Gentzen’s sequent calculus proof rules is the construction of (possibly) large *synthetic inference rules*. In this paper, we examine the synthetic inference rules that arise when using theories composed of bipolars, and we do this in both classical and intuitionistic logics. A key step in transforming a formula into synthetic inference rules involves attaching a *polarity* to atomic formulas and to some logical connectives. Since there are different choices in how polarity is assigned, it is possible to produce different synthetic inference rules for the same formula. We show that this flexibility allows for the generalization of different approaches for transforming axioms into sequent rules present in the literature. We finish the paper showing how to apply these results to organize the proof theory of labeled sequent systems for several propositional modal logics.

1. Introduction

We start by presenting a simple, motivating example that should illustrate several key concepts we shall encounter. Let B be a formula and Γ be a multiset of formulas. Consider attempting to build a proof of the following two-sided sequent

$$\Gamma, A_1 \supset \cdots \supset A_n \supset A_0 \vdash B,$$

in which the distinguished implication is such that $n \geq 1$ and A_0, \dots, A_n are atomic formulas. In general, there are many ways to proceed with attempting to build a cut-free proof of this sequent and we characterize them as one of the following four possibilities. This sequent can be the conclusion of

1. a structural rule (weakening or contraction) or the initial rule;
2. a right introduction rule, if B is not an atomic formula;
3. a left-introduction rule that introduces a formula in Γ ; or
4. the implication-introduction rule that introduces the distinguished implication.

The number of possible choices here could be large, particularly if Γ contains a large number of formulas. If we chose the fourth of these possibilities, the proof would look as follows (at least in the intuitionistic setting):

$$\frac{\Gamma \vdash A_1 \quad \Gamma, A_2 \supset \cdots \supset A_n \supset A_0 \vdash B}{\Gamma, A_1 \supset \cdots \supset A_n \supset A_0 \vdash B} L\supset$$

Note that we again have a large number of possible ways to proceed in attempting to prove the right premise: indeed, if $n \geq 2$, we have all the same choices as before. Clearly, those choices—and their multiplicative effects as we search for a sequence of inference steps that terminates in a proof—are in desperate need of being structured somehow. Focused proof systems provide such structure using the following two devices.

Focused rule application If you chose to apply the implication-left introduction on the distinguished implication, then you also commit to repeat the implication-left rule on the right premise until the atomic formula A_0 results. That is, the left-introduction applied to the distinguished implication results in the following derived inference rule

$$\frac{\Gamma \vdash A_1 \quad \dots \quad \Gamma \vdash A_n \quad \Gamma, A_0 \vdash B}{\Gamma, A_1 \supset \cdots \supset A_n \supset A_0 \vdash B} L\supset \quad n \text{ times.}$$

Polarization Although the focused application of inference rules provides structure to attempts to build proofs, there are still so many remaining choices, that it is possible to impose two different “protocols” for restricting choices further. The Q -protocol insists that the first n premises above are trivial, meaning that they are proved by the initial rule. Following that protocol, we have $A_i \in \Gamma$ for $1 \leq i \leq n$. Thus, if we set Γ' to be the result of removing all occurrences of A_1, \dots, A_n from Γ , then the derived inference rule above becomes

$$\frac{\Gamma', A_1, \dots, A_n, A_0 \vdash B}{\Gamma', A_1, \dots, A_n, A_1 \supset \cdots \supset A_n \supset A_0 \vdash B} .$$

The second protocol, the T -protocol insists that the right-most premise is trivial: that is, A_0 and B are the same atomic formula. Thus, the derived inference rule above becomes

$$\frac{\Gamma \vdash A_1 \quad \dots \quad \Gamma \vdash A_n}{\Gamma, A_1 \supset \cdots \supset A_n \supset A_0 \vdash A_0} .$$

Using the Q -protocol, the proof-search semantics of the implication $A_1 \supset \dots \supset A_n \supset A_0$ is given by *forward-chaining*: if you have assumptions A_1, \dots, A_n then you can add the assumption A_0 . Using the T -protocol, the proof-search semantics of the same implication is given by *back-chaining*: in order to prove the conclusion A_0 , attempt instead to prove each of A_1, \dots, A_n . The names for the Q and T protocols comes from Danos, Joinet, and Schellinx [1]: in the Q protocol, the tail (“queue”) of an implication yields a trivial premise while in the T protocol, the head (“tête”) of an implication yields a trivial premise. A more modern and flexible presentation of the Q and T protocols speaks, instead, of the *polarity* of formulas: for this example, the polarity given to atomic formulas is the most relevant. In particular, if all atomic formulas have a positive polarity, the Q -protocol is enforced, while if all atomic formulas have a negative polarity, the T -protocol is enforced.

In the next section, we introduce the *LKF* and *LJF* [2, 3] focused proof systems for classical and intuitionistic logics, respectively. Those systems extend both the notion of focusing and polarity to all formulas, moving beyond the example above involving only implications and atomic formulas. In particular, *focused rule applications* imply that focus is transferred from conclusion to premises in derivations. This process goes on until either the focused phase ends (depending on the *polarity* of the focused formula), or the derivation ends. Once the focus is released, the formula is eagerly decomposed into subformulas, which are ultimately *stored* in the context.

Reading derivations from the root upwards, this forces a sequent derivation to be organized into *focused phases*, each of them corresponding to an application of a *synthetic inference rule* [4], where the focused formula is rewritten into (some of) its subformulas.

There is a class of formulas corresponding to particularly interesting synthetic rules: the *bipolars* (Section 3). Bipolars are formulas in which polarity can change at most once among its subformulas. This means that focusing on a bipolar A gives rise to (possibly many) synthetic inference rules having simple shape, with leaves involving only atomic subformulas of A . We call a synthetic inference rule corresponding to the bipolar A a *bipole* for A .

In this work, we will present a careful study of bipoles, giving a fresh view to an old problem: how to incorporate inference rules encoding axioms into proof systems for classical and intuitionistic logics (Section 4).

A key step in transforming a formula into synthetic inference rules involves attaching a polarity to atomic formulas and to some logical connectives. Since there are different choices for assigning polarities, it is possible to produce different synthetic inference rules for the same (unpolarized) formula. In the example above, there are (at most) 2^{n+1} different possible polarizations for the atomic formulas in $A_1 \supset \dots \supset A_n \supset A_0$, each of them corresponding to a different bipole. We show that this flexibility allows for the generalization of different approaches for transforming axioms into sequent rules present in the literature.

The general problem of extending standard proof-theoretical results obtained for pure logic to certain class of non-logical axioms has been focus of attention for

quite some time now. The main challenge in this effort is to determine a general procedure that guarantees that such extensions preserve good proof-theoretical properties.

A remarkable step in that direction was the careful investigation of *geometric axioms*. Geometric axioms are first-order formulas that can be converted into (natural deduction/sequent) inference rules having “a certain simple form in which only atomic formulas play a critical part”, as described by Simpson [5]. And this “simple rules for atomic formulas” motto seems to be the core of success in this endurance in the approaches/extensions present in the literature [6].

The fact that bipolars smoothly extend geometric formulas was first noted in [7] where, among other things, Ciabattoni et al. developed a systematic procedure for transforming a class of (Hilbert) axioms (called \mathcal{N}_2) into equivalent *structural* inference rules in substructural sequent calculi. Extended to first-order logic [8], such a procedure showed how to formalize and generalize the results in [9, 10, 11], where Negri et al. proposed methods for transforming axioms into inference rules, without affecting the admissibility of the structural rules. Bipolars are actually first-order atomic-polarized versions of \mathcal{N}_2 formulas.

Following a parallel path, Viganò presents in [12] a detailed study of extensions of modal systems with Horn relational theories – a sub-class of geometric theories. Interestingly enough, the difference between Viganò and Negri’s approaches is only the *protocol*: back-chaining *versus* forward-chaining, respectively.

In this work, we come back to the inception of the axioms-as-rules problem, showing that the combination of bipolars and focusing is the real essence of “simple rules for atomic formulas”. This implies that the previously mentioned works are different faces of the same coin, minted from focusing and polarization. Moreover, we address these issues with a uniform presentation in both classical and intuitionistic first-order logics.

We finish the paper by showing how to emulate precisely rules for modalities in labeled modal systems as synthetic connectives [13, 14] (Section 5). Such tight emulation means that proof search and proof checking on the focused version of the translated formulas imitates exactly proof search and proof checking in the correspondent labeled system. As a result, we are able to show that we can use focused proofs to precisely emulate modal proofs whenever Kripke frames are characterized by bipolar properties.

2. Background notions

The formulas of first-order classical and intuitionistic logics are built from atomic formula along with t and f (true and false), \wedge and \vee (conjunction and disjunction), \supset (implication), and \forall , \exists (universal and existential quantification). The *logical units* are t and f , the *binary connectives* are \wedge , \vee , and \supset , and the *quantifiers* are \forall and \exists . Collectively, these are all called *logical connectives*. Here, we do not take negation as a logical connective, instead we write $B \supset f$ to denote the negation of B .

INTRODUCTION RULES

$$\begin{array}{c}
\frac{A, \Gamma \vdash \Delta}{A \wedge B, \Gamma \vdash \Delta} \quad \frac{B, \Gamma \vdash \Delta}{A \wedge B, \Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta, A \quad \Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \wedge B} \quad \frac{}{\Gamma \vdash \Delta, t} \\
\\
\frac{A, \Gamma \vdash \Delta \quad B, \Gamma \vdash \Delta}{A \vee B, \Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta, A}{\Gamma \vdash \Delta, A \vee B} \quad \frac{\Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \vee B} \quad \frac{}{f, \Gamma \vdash \Delta} \\
\\
\frac{\Gamma \vdash \Delta_1, A \quad B, \Gamma \vdash \Delta_2}{A \supset B, \Gamma \vdash \Delta_1, \Delta_2} \quad \frac{A, \Gamma \vdash \Delta, B}{\Gamma \vdash \Delta, A \supset B} \\
\\
\frac{[t/x]B, \Gamma \vdash \Delta}{\forall x B, \Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta, [y/x]B}{\Gamma \vdash \Delta, \forall x B} \quad \frac{[y/x]B, \Gamma \vdash \Delta}{\exists x B, \Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta, [t/x]B}{\Gamma \vdash \Delta, \exists x B}
\end{array}$$

IDENTITY RULE (the initial rule)

$$\frac{}{P, \Gamma \vdash \Delta, P}$$

STRUCTURAL RULES (the contraction rules)

$$\frac{A, A, \Gamma \vdash \Delta}{A, \Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta, A, A}{\Gamma \vdash \Delta, A}$$

Figure 1: The classical sequent calculus LK . The LJ calculus results from restricting the right-hand side of sequents to contain at most one formula. Here, A and B are arbitrary formulas, and P is an atomic formula. In the \forall right rule and in the \exists left rule, the eigenvariable y does not occur free in any formula of the conclusion.

2.1. Sequent calculus proof systems for classical and intuitionistic logic

Figure 1 contains the sequent calculus inference rules for what we shall call LK . This system is formally different from the one of the same name given by Gentzen in [15]. In particular, Gentzen’s original system included the cut inference rule, but we delay until Section 3 to introduce that inference rule, and when we do it will be as an inference rule in a focused proof system. Other differences from Gentzen’s original LK proof system—such as the restriction of the initial rule to atomic formulas—do not change the character of the LK proof system in any important fashion. Just as in [15], the intuitionistic system LJ is obtained by restricting the right side of each sequent to contain at most one formula. This restriction on the right-hand side of sequents is equivalent to the following two restrictions: (i) no contractions are allowed on the right side and (ii) the rule for introducing implication on the left side is restricted to have the

form

$$\frac{\Gamma \vdash A \quad B, \Gamma \vdash \Delta}{A \supset B, \Gamma \vdash \Delta}.$$

That is, the right-hand side of the conclusion must be the same as the right-hand side of the right premise.

2.2. Polarized formulas

An early focused proof, given in [16, 17], introduced a two-phase proof system—used to capture the logic programming concepts of *goal-reduction* and *back-chaining*—and proved it to be complete for a subset of intuitionistic logic. In [18], Andreoli generalized that two phase construction by extending it to all of linear logic. Subsequently, several additional proof systems appeared in which somewhat similar proof structures were given for classical and intuitionistic logics: in particular, *LKT* and *LKQ* [1], *LJT* [19], and *LJQ* [20]. The focused proof systems *LKF* and *LJF* [2, 3] were designed to generalize all of those proof systems: in particular, *LKF* and *LJF* can accommodate both the *Q* and *T* protocols as well as a mix of those protocols. The proof system *LKF*, for first-order classical logic, and the proof system *LJF*, for first-order intuitionistic logic, are presented in Figures 2 and 3, respectively. Our presentation has been adapted from the corresponding proof systems given in [3]: in particular, for ease of comparison between the intuitionistic and the classical proofs, the proof system *LKF* is presented using two-sided sequents.

In order to obtain their flexibility in capturing various focusing regimes, the *LKF* and *LJF* proof systems use *polarized* formulas instead of the unpolarized formulas used in the *LK* and *LJ* proof systems of Section 2.1. A *polarized classical (first-order) formula* is a formula built using atomic formulas, the usual first-order quantifiers \forall and \exists , the implication \supset , and polarized versions of the logical connectives and constants, i.e., t^- , t^+ , f^- , f^+ , \vee^- , \vee^+ , \wedge^- , \wedge^+ . A *polarized intuitionistic (first-order) formula* is a polarized classical formula in which the logical connectives f^- and \vee^- do not occur. The positive and negative versions of connectives and constants have identical truth conditions but different inference rules inside the polarized proof systems. For example, the left introduction rule for \wedge^+ is invertible while the left introduction rule for \wedge^- is not invertible.

We shall also find it necessary to use *delays*: if B is a polarized formula then we define $\partial_-(B)$ to be (the always negative) $B \wedge^- t^-$ and $\partial_+(B)$ to be (the always positive) $B \wedge^+ t^+$. Equivalently, we can take $\partial_+(\cdot)$ to be the 1-ary version of either the binary \vee^+ or \wedge^+ and take $\partial_-(\cdot)$ to be the 1-ary version of either the binary \vee^- or \wedge^- . (The 0-ary version of these four connectives correspond to the logical units f^+ , t^+ , f^- , t^- .)

If a formula's top-level connective is t^+ , f^+ , \vee^+ , \wedge^+ , or \exists , then that formula is *positive*. If a formula's top-level connective is t^- , f^- , \vee^- , \wedge^- , \supset , or \forall , then it is *negative*. Note that in the intuitionistic system *LJF*, we have only one disjunction and one falsum, both of which exist only with positive polarity. The way to form the negation of the polarized formula B is with the formula $B \supset f^+$: this formula has negative polarity no matter the polarity of B .

In both *LKF* and *LJF*, every polarized formula is classified as positive or negative. This means that we must also provide a polarity to atomic formulas. As it turns out, this assignment of polarity to atomic formulas can, in principle, be arbitrary. In particular, an *atomic bias assignment* is a function $\delta(\cdot)$ that maps atomic formulas to the set of two tokens $\{pos, neg\}$: if $\delta(A)$ is *pos* then that atomic formula is positive and if $\delta(A)$ is *neg* then that atomic formula is negative. We may ask that all atomic formulas are positive, that they are all negative, or we can mix polarity assignments. In particular, the atomic bias assignment $\delta^+(\cdot)$ assigns all atoms a positive polarity while $\delta^-(\cdot)$ assigns all atoms a negative polarity. For this paper, we shall assume that an atomic bias assignment is also *stable under substitution*: that is, for all substitutions θ , $\delta(\theta A) = \delta(A)$. In first-order logic, this is equivalent to saying that such bias assignments are *predicate determined*: that is, if atoms A and A' have the same predicate head, then $\delta(A) = \delta(A')$.

We say that the pair $\langle \delta, \hat{B} \rangle$ is a *polarization of B* if $\delta(\cdot)$ is an atomic bias assignment and if every occurrence of t , \wedge , f , and \vee in B is labeled with either the $+$ or $-$ annotation. If B has n occurrences of these logical connectives then there are 2^n different ways to place these $+$ or $-$ symbols. We shall also allow the insertion of any number of $\partial_+(\cdot)$ and $\partial_-(\cdot)$ into \hat{B} as well. In other words, the polarized formula $\langle \delta, C \rangle$ is a polarization of B if deleting all delays and all $+$ and $-$ annotations on logical connectives of C results in B . Note that we use \supset , \forall , and \exists in both unpolarized as well as polarized formulas: we can do this since the polarity of these connectives is not ambiguous. In classical logic, the polarity of t , \wedge , f , and \vee is ambiguous and all of these can be positive or negative. In intuitionistic logic, only the polarity of t and \wedge is ambiguous. In both of these logics, however, the polarity of atoms is equally ambiguous. Finally, if $\langle \delta, \hat{B} \rangle$ is a polarization of B , we shall generally drop explicit reference to δ and simply say that \hat{B} is a polarization of B : often, the atomic bias assignment is either not important or can be inferred from context.

2.3. Focused proof systems

The inference rules of *LKF* and *LJF* presented in Figures 2 and 3, respectively, involve three kinds of sequents

$$\Gamma \uparrow \Theta \vdash \Omega \uparrow \Delta, \quad \Gamma \Downarrow B \vdash \Delta, \quad \text{and} \quad \Gamma \vdash B \Downarrow \Delta,$$

where Γ , Θ , Ω and Δ are multisets of polarized formulas and B is a polarized formula. The formula occurrence B in a \Downarrow -sequent is called the *focus* of that sequent.

The system *LJF* is depicted in a separate figure for the sake of clarity. However, one can notice that, similarly to what we have for *LJ* and *LK* in the original Gentzen formulations, *LJF* can be seen as a restriction of *LKF*, where the rules for f^- and \vee^- are omitted and only one formula is admitted in the succedent of sequents. In particular, this implies that (i) in the left rule for \supset , the right context of the conclusion is not present in the left premise; (ii) in the rule D_r , the formula placed under focus is not subjected to contraction; and (iii)

ASYNCHRONOUS RULES

$$\begin{array}{c}
\frac{\Gamma \uparrow \Theta \vdash A, B, \Omega \uparrow \Delta}{\Gamma \uparrow \Theta \vdash A \vee \neg B, \Omega \uparrow \Delta} \vee_r^- \quad \frac{\Gamma \uparrow \Theta \vdash A, \Omega \uparrow \Delta \quad \Gamma \uparrow \Theta \vdash B, \Omega \uparrow \Delta}{\Gamma \uparrow \Theta \vdash A \wedge \neg B, \Omega \uparrow \Delta} \wedge_r^- \\
\frac{\Gamma \uparrow A, B, \Theta \vdash \Omega \uparrow \Delta}{\Gamma \uparrow A \wedge^+ B, \Theta \vdash \Omega \uparrow \Delta} \wedge_l^+ \quad \frac{\Gamma \uparrow A, \Theta \vdash \Omega \uparrow \Delta \quad \Gamma \uparrow B, \Theta \vdash \Omega \uparrow \Delta}{\Gamma \uparrow A \vee^+ B, \Theta \vdash \Omega \uparrow \Delta} \vee_l^+ \\
\frac{\Gamma \uparrow \cdot \vdash [y/x]B, \Omega \uparrow \Delta}{\Gamma \uparrow \cdot \vdash \forall x.B, \Omega \uparrow \Delta} \forall_r \quad \frac{\Gamma \uparrow [y/x]B, \Theta \vdash \Omega \uparrow \Delta}{\Gamma \uparrow \exists x.B, \Theta \vdash \Omega \uparrow \Delta} \exists_l \\
\frac{\Gamma \uparrow \Theta, A \vdash B, \Omega \uparrow \Delta}{\Gamma \uparrow \Theta \vdash A \supset B, \Omega \uparrow \Delta} \supset_r \quad \frac{\Gamma \uparrow \Theta \vdash \Omega \uparrow \Delta}{\Gamma \uparrow t^+, \Theta \vdash \Omega \uparrow \Delta} t_l^+ \quad \frac{\Gamma \uparrow \Theta \vdash \Omega \uparrow \Delta}{\Gamma \uparrow \Theta \vdash f^-, \Omega \uparrow \Delta} f_r^- \\
\frac{}{\Gamma \uparrow \Theta \vdash t^-, \Omega \uparrow \Delta} t_r^- \quad \frac{}{\Gamma \uparrow f^+, \Theta \vdash \Omega \uparrow \Delta} f_l^+
\end{array}$$

SYNCHRONOUS RULES

$$\begin{array}{c}
\frac{\Gamma \vdash A \Downarrow \Delta \quad \Gamma \Downarrow B \vdash \Delta}{\Gamma \Downarrow A \supset B \vdash \Delta} \supset_l \quad \frac{\Gamma \Downarrow A \vdash \Delta \quad \Gamma \Downarrow B \vdash \Delta}{\Gamma \Downarrow A \vee \neg B \vdash \Delta} \vee_l^- \quad \frac{\Gamma \Downarrow A_i \vdash \Delta}{\Gamma \Downarrow A_1 \wedge \neg A_2 \vdash \Delta} \wedge_l^- \\
\frac{\Gamma \vdash A \Downarrow \Delta \quad \Gamma \vdash B \Downarrow \Delta}{\Gamma \vdash A \wedge^+ B \Downarrow \Delta} \wedge_r^+ \quad \frac{\Gamma \vdash A_i \Downarrow \Delta}{\Gamma \vdash A_1 \vee^+ A_2 \Downarrow \Delta} \vee_r^+ \\
\frac{\Gamma \Downarrow [t/x]B \vdash \Delta}{\Gamma \Downarrow \forall x.B \vdash \Delta} \forall_l \quad \frac{\Gamma \vdash [t/x]B \Downarrow \Delta}{\Gamma \vdash \exists x.B \Downarrow \Delta} \exists_r \quad \frac{}{\Gamma \vdash t^+ \Downarrow \Delta} t_r^+ \quad \frac{}{\Gamma \Downarrow f^- \vdash \Delta} f_l^-
\end{array}$$

IDENTITY RULES

$$\frac{}{\Gamma \Downarrow N_a \vdash N_a, \Delta} l_l \quad \frac{}{\Gamma, P_a \vdash P_a \Downarrow \Delta} l_r$$

STRUCTURAL RULES

$$\begin{array}{c}
\frac{\Gamma, N \Downarrow N \vdash \Delta}{\Gamma, N \uparrow \cdot \vdash \cdot \uparrow \Delta} D_l \quad \frac{\Gamma \vdash P \Downarrow P, \Delta}{\Gamma \uparrow \cdot \vdash \cdot \uparrow P, \Delta} D_r \quad \frac{\Gamma \uparrow P \vdash \cdot \uparrow \Delta}{\Gamma \Downarrow P \vdash \Delta} R_l \quad \frac{\Gamma \uparrow \cdot \vdash N \uparrow \Delta}{\Gamma \vdash N \Downarrow \Delta} R_r \\
\frac{C, \Gamma \uparrow \Theta \vdash \Omega \uparrow \Delta}{\Gamma \uparrow C, \Theta \vdash \Omega \uparrow \Delta} S_l \quad \frac{\Gamma \uparrow \cdot \vdash \Omega \uparrow D, \Delta}{\Gamma \uparrow \cdot \vdash D, \Omega \uparrow \Delta} S_r
\end{array}$$

Here, P is positive, N is negative, C is a negative formula or positive atom, D a positive formula or negative atom, N_a is a negative atom, and P_a is a positive atom. Other formulas are arbitrary. In the rules \forall_r and \exists_l the eigenvariable y does not occur free in any formula of the conclusion.

Figure 2: The focused classical sequent calculus *LKF*.

ASYNCHRONOUS RULES

$$\begin{array}{c}
\frac{\Gamma \uparrow A, \Theta \vdash B \uparrow \cdot}{\Gamma \uparrow \Theta \vdash A \supset B \uparrow \cdot} \supset_r \quad \frac{\Gamma \uparrow \Theta \vdash A \uparrow \cdot \quad \Gamma \uparrow \Theta \vdash B \uparrow \cdot}{\Gamma \uparrow \Theta \vdash A \wedge B \uparrow \cdot} \wedge_r \\
\frac{\Gamma \uparrow A, B, \Theta \vdash \mathcal{R}}{\Gamma \uparrow A \wedge^+ B, \Theta \vdash \mathcal{R}} \wedge_l^+ \quad \frac{\Gamma \uparrow A, \Theta \vdash \mathcal{R} \quad \Gamma \uparrow B, \Theta \vdash \mathcal{R}}{\Gamma \uparrow A \vee^+ B, \Theta \vdash \mathcal{R}} \vee_l^+ \\
\frac{\Gamma \uparrow \Theta \vdash [y/x]B \uparrow \cdot}{\Gamma \uparrow \Theta \vdash \forall x.B \uparrow \cdot} \forall_r \quad \frac{\Gamma \uparrow [y/x]B, \Theta \vdash \mathcal{R}}{\Gamma \uparrow \exists x.B, \Theta \vdash \mathcal{R}} \exists_l \\
\frac{}{\Gamma \uparrow \Theta \vdash t^- \uparrow \cdot} t_r^- \quad \frac{\Gamma \uparrow \Theta \vdash \mathcal{R}}{\Gamma \uparrow t^+, \Theta \vdash \mathcal{R}} t_l^+ \quad \frac{}{\Gamma \uparrow f^+, \Theta \vdash \mathcal{R}} f_l^+
\end{array}$$

SYNCHRONOUS RULES

$$\begin{array}{c}
\frac{\Gamma \vdash A \downarrow \quad \Gamma \downarrow B \vdash R}{\Gamma \downarrow A \supset B \vdash R} \supset_l \quad \frac{\Gamma \vdash A_i \downarrow}{\Gamma \vdash A_1 \vee^+ A_2 \downarrow} \vee_r^+ \quad \frac{\Gamma \downarrow A_i \vdash R}{\Gamma \downarrow A_1 \wedge^- A_2 \vdash R} \wedge_l^- \\
\frac{\Gamma \vdash A \downarrow \quad \Gamma \vdash B \downarrow}{\Gamma \vdash A \wedge^+ B \downarrow} \wedge_r^+ \quad \frac{\Gamma \downarrow [t/x]B \vdash R}{\Gamma \downarrow \forall x.B \vdash R} \forall_l \quad \frac{\Gamma \vdash [t/x]B \downarrow}{\Gamma \vdash \exists x.B \downarrow} \exists_r \quad \frac{}{\Gamma \vdash t^+ \downarrow} t_r^+
\end{array}$$

IDENTITY RULES

$$\frac{}{\Gamma \downarrow N_a \vdash N_a} l_l \quad \frac{}{\Gamma, P_a \vdash P_a \downarrow} l_r$$

STRUCTURAL RULES

$$\begin{array}{c}
\frac{\Gamma, N \downarrow N \vdash R}{\Gamma, N \uparrow \cdot \vdash \cdot \uparrow R} D_l \quad \frac{\Gamma \vdash P \downarrow}{\Gamma \uparrow \cdot \vdash \cdot \uparrow P} D_r \quad \frac{\Gamma \uparrow P \vdash \cdot \uparrow R}{\Gamma \downarrow P \vdash R} R_l \quad \frac{\Gamma \uparrow \cdot \vdash N \uparrow \cdot}{\Gamma \vdash N \downarrow} R_r \\
\frac{C, \Gamma \uparrow \Theta \vdash \mathcal{R}}{\Gamma \uparrow C, \Theta \vdash \mathcal{R}} S_l \quad \frac{\Gamma \uparrow \cdot \vdash \cdot \uparrow D}{\Gamma \uparrow \cdot \vdash D \uparrow \cdot} S_r
\end{array}$$

Here, P is positive, N is negative, C is a negative formula or positive atom, D a positive formula or negative atom, N_a is a negative atom, and P_a is a positive atom. Other formulas are arbitrary. \mathcal{R} denotes $\Delta_1 \uparrow \Delta_2$ where the union of Δ_1 and Δ_2 contains at most one formula. In the rules \forall_r and \exists_l the eigenvariable y does not occur free in any formula of the conclusion.

Figure 3: The focused intuitionistic sequent calculus *LJF*.

a sequent of the form $\Gamma \vdash B \Downarrow \Delta$, when used in an *LJF* proof, is such that Δ is empty. In that case, we write that sequent as simply $\Gamma \vdash B \Downarrow$.

The soundness of *LKF* and *LJF* can be stated as follows.

1. Let B be an unpolarized classical logic formula and let \hat{B} be any polarization of B . If $\cdot \Uparrow \cdot \vdash \hat{B} \Uparrow \cdot$ is provable in *LKF* then $\vdash B$ is provable in *LK*.
2. Let B be an unpolarized intuitionistic logic formula and let \hat{B} be any polarization of B . If $\cdot \Uparrow \cdot \vdash \hat{B} \Uparrow \cdot$ is provable in *LJF* then $\vdash B$ is provable in *LJ*.

These claims of soundness will be proved in detail in Section 2.4.

The completeness of *LKF* and *LJF* can be stated as follows.

Theorem 1 (Completeness of *LKF* and *LJF*). 1. *If B is an unpolarized classical logic theorem (i.e., $\vdash B$ is provable in *LK*) and \hat{B} is any polarization of B , then $\cdot \Uparrow \cdot \vdash \hat{B} \Uparrow \cdot$ is provable in *LKF*.*
 2. *If B is an unpolarized intuitionistic logic theorem (i.e., $\vdash B$ is provable in *LJ*) and \hat{B} is any polarization of B , then $\cdot \Uparrow \cdot \vdash \hat{B} \Uparrow \cdot$ is provable in *LJF*.*

The proofs of these completeness theorems are lengthy and are not given here: the interested reader can find them in [3, 21]. A consequence of the completeness theorem is that the choice of polarization does not affect provability (although it can have an impact on the structure of proofs). Hence, if a polarization of B is provable in *LKF* (*LJF*) then every polarization of B is provable in *LKF* (respectively, *LJF*).

We shall now make an important distinction between the terms *derivation* and *proof*. While they are both tree-structured organizations of inference rules (focused or not), we shall only use the term “proof” when all leaves of that tree are closed: that is, all leaves are justified by either an initial rule (l_l, l_r) or the introduction of a logical unit ($t_r^-, t_r^+, f_l^-, f_l^+$). Derivations can have zero or more leaves that are not the consequence of an inference rule.

By observing *LKF* and *LJF* inference rules in Figures 2 and 3, we notice that derivations are constructed by a repeated alternation of two phases: a *synchronous phase*, which (reading the derivation from the root upwards) typically starts with the application of a *decide* rule (D_l, D_r) and consists in the application of synchronous rules, and an *asynchronous phase*, which starts with the application of a *release* rule (R_l, R_r), and consists in the application of asynchronous rules, terminating with applications of a *store* rule (S_l, S_r).

Definition 2 (Synthetic inference rule). *A synthetic inference rule is an inference rule of the form*

$$\frac{\Gamma_1 \Uparrow \cdot \vdash \cdot \Uparrow \Delta_1 \quad \dots \quad \Gamma_n \Uparrow \cdot \vdash \cdot \Uparrow \Delta_n}{\Gamma \Uparrow \cdot \vdash \cdot \Uparrow \Delta}$$

which is justified by a derivation of the form

$$\frac{\Gamma_1 \Uparrow \cdot \vdash \cdot \Uparrow \Delta_1 \quad \dots \quad \Gamma_n \Uparrow \cdot \vdash \cdot \Uparrow \Delta_n}{\Gamma \Uparrow \cdot \vdash \cdot \Uparrow \Delta} \Pi$$

Here, $n \geq 0$ and the inference rules of derivation Π are such that no synchronous rule application occurs above an asynchronous rule application. We also assume that Π contains at least one inference rule.

Sequents of the form $\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta$ are called *border sequents* since they form the borders (the endsequent and premises) of synthetic inference rules. We will occasionally identify a synthetic inference rule with the derivation justifying it. We can speak of such synthetic inference rules in both *LKF* and *LJF* and, in both cases, the last inference rule of (the justification) Π must be a decide rule, either D_l or D_r . In the case that that decide rule is D_l and it selects for focus the (negative) formula B , we say that this derivation is a *synthetic inference rule for B* .

Our main use of focused proofs in this paper is to examine synthetic inference rules for formulas from certain theories.

2.4. Encoding unfocused systems in focused systems

In the introduction, we motivated using focusing and polarization to build large-scale inference rules, such as forward-chaining or back-chaining. In this section, we start with showing that the small-scale introduction rules in Gentzen's original, unfocused proof systems can be emulated precisely using synthetic inference rules in the corresponding focused proof system. Once we develop the techniques to show that emulation, we will move to addressing larger-scale inference rules in Section 3.

Given a polarized formula B we will denote by B° the first-order formula obtained by removing the annotations on the polarized versions of conjunction, disjunction, true, and false. For example, if P , Q , and R are atoms, then

$$(P \vee^- (Q \wedge^+ R))^\circ = P \vee (Q \wedge R)$$

This translation carries \forall , \exists , \supset , and atoms to themselves. If Γ is a set of polarized formulas, then Γ° denotes the set $\{B^\circ \mid B \in \Gamma\}$.

It is straightforward to transform a derivation Π of $\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta$ in *LKF* (or *LJF*) into a derivation Π° of $\Gamma^\circ \vdash \Delta^\circ$ in *LK* (respectively, *LJ*) by ignoring the release and store rules and by using the contraction rule when replacing the decide rules and when transforming multiplicative rules into additive ones. For example, if P, Q, R are atomic formulas assigned positive polarity, the *LKF* derivation

$$\frac{\frac{\frac{P \vee^- (Q \wedge^+ R), P \uparrow \cdot \vdash \cdot \uparrow \cdot}{P \vee^- (Q \wedge^+ R) \uparrow P \vdash \cdot \uparrow \cdot} S_l \quad \frac{\frac{P \vee^- (Q \wedge^+ R), Q, R \uparrow \cdot \vdash \cdot \uparrow \cdot}{P \vee^- (Q \wedge^+ R) \uparrow Q, R \vdash \cdot \uparrow \cdot} S_l}{P \vee^- (Q \wedge^+ R) \uparrow Q \wedge^+ R \vdash \cdot \uparrow \cdot} \wedge_l^+ \quad \frac{\frac{P \vee^- (Q \wedge^+ R) \uparrow P \vdash \cdot \uparrow \cdot}{P \vee^- (Q \wedge^+ R) \downarrow P \vdash \cdot} R_l \quad \frac{\frac{P \vee^- (Q \wedge^+ R), Q, R \uparrow \cdot \vdash \cdot \uparrow \cdot}{P \vee^- (Q \wedge^+ R) \uparrow Q, R \vdash \cdot \uparrow \cdot} S_l}{P \vee^- (Q \wedge^+ R) \uparrow Q \wedge^+ R \vdash \cdot \uparrow \cdot} \wedge_l^+}{P \vee^- (Q \wedge^+ R) \downarrow Q \wedge^+ R \vdash \cdot} R_l}{P \vee^- (Q \wedge^+ R) \downarrow P \vee^- (Q \wedge^+ R) \vdash \cdot} \vee_l}{\frac{P \vee^- (Q \wedge^+ R) \downarrow P \vee^- (Q \wedge^+ R) \vdash \cdot}{P \vee^- (Q \wedge^+ R) \uparrow \cdot \vdash \cdot \uparrow \cdot} D_l} \vee_l$$

is transformed into the *LK* derivation

$$\frac{\frac{\frac{P \vee (Q \wedge R), Q, R \vdash}{P \vee (Q \wedge R), Q, Q \wedge R \vdash}}{P \vee (Q \wedge R), Q \wedge R, Q \wedge R \vdash}}{P \vee (Q \wedge R), P \vdash} \frac{P \vee (Q \wedge R), Q \wedge R \vdash}{P \vee (Q \wedge R), Q \wedge R \vdash}}{P \vee (Q \wedge R), P \vee (Q \wedge R) \vdash} \frac{P \vee (Q \wedge R) \vdash}{P \vee (Q \wedge R) \vdash}$$

It is possible to map unfocused *LK* proofs into *LKF* proofs in such a way that every rule application in *LK* corresponds to a synthetic inference rule in *LKF*. To do such an emulation, it is important to break up a sequence of negative or positive connectives, by inserting the delays $\partial_+(\cdot)$ and $\partial_-(\cdot)$. From the definitions given for delays in Section 2.2, the following additional focused inference rules can be justified.

$$\frac{\Gamma \vdash B \Downarrow \Delta}{\Gamma \vdash \partial_+(B) \Downarrow \Delta} \partial_+^r \quad \frac{\Gamma \Uparrow \Theta, B \vdash \Omega \Uparrow \Delta}{\Gamma \Uparrow \Theta, \partial_+(B) \vdash \Omega \Uparrow \Delta} \partial_+^l$$

$$\frac{\Gamma \Uparrow \Theta \vdash B, \Omega \Uparrow \Delta}{\Gamma \Uparrow \Theta \vdash \partial_-(B), \Omega \Uparrow \Delta} \partial_-^r \quad \frac{\Gamma \Downarrow B \vdash \Delta}{\Gamma \Downarrow \partial_-(B) \vdash \Delta} \partial_-^l$$

Following [3], we define the *left/right* translation functions $[\cdot]^l$ and $[\cdot]^r$ from first-order formulas into polarized formulas recursively as follows: if P is an atom, then $[P]^l = [P]^r = P$; otherwise

$$\begin{aligned} [f]^l &= \partial_-(f^+) & [t]^l &= \partial_-(t^+) \\ [A \wedge B]^l &= \partial_+([A]^l) \wedge^- \partial_+([B]^l) & [A \vee B]^l &= \partial_-([A]^l) \vee^+ [B]^l \\ [A \supset B]^l &= \partial_-([A]^r) \supset \partial_+([B]^l) & [\forall x.A]^l &= \forall x.\partial_+([A]^l) \\ [\exists x.A]^l &= \partial_-([\exists x.A]^l) \\ [f]^r &= f^+ & [t]^r &= \partial_+(t^-) \\ [A \wedge B]^r &= \partial_+([A]^r) \wedge^- [B]^r) & [A \vee B]^r &= \partial_-([A]^r) \vee^+ \partial_-([B]^r) \\ [A \supset B]^r &= \partial_+([A]^l \supset [B]^r) & [\forall x.A]^r &= \partial_+(\forall x.[A]^r) \\ [\exists x.A]^r &= \exists x.\partial_-([A]^r) \end{aligned}$$

Since these translations do not use either f^- or \vee^- , the resulting formulas can be used in both the *LJF* and *LKF* proof systems. These translations do not assign any polarity to atomic formulas and any atomic polarity assignment can be paired with the result of such translations. The translations are applied to multisets, say Ω , of polarized formulas in the usual way: $[\Omega]^r = \{[A]^r \mid A \in \Omega\}$ and $[\Omega]^l = \{[A]^l \mid A \in \Omega\}$. Finally, we define a translation from *LK* (*LJ*) sequents into *LKF* (*LJF*) border sequents as follows:

$$[(A_1, \dots, A_n \vdash B_1, \dots, B_m)] = [A_1, \dots, A_n]^l \Uparrow \cdot \vdash \cdot \Uparrow [B_1, \dots, B_m]^r.$$

In both translation of multisets of formulas and of sequents, we shall assume that only one atomic polarity assignment is used for all formulas in these collections of polarized formulas.

Note that if B is a non-atomic unpolarized formula then $[B]^l$ is always negative while $[B]^r$ is always positive. If B is atomic, then the polarity of $[B]^r$ and $[B]^l$ is the same as the polarity given by the atomic polarity assignment.

As we show below, the delays in $[\cdot]^l$ and $[\cdot]^r$ break focusing phases and this allows us to mimic the small step inference rules of LK and LJ derivations in LKF and LJF , respectively (see also [3]).

2.4.1. Mapping unpolarized proofs to polarized proofs

One important difference between the unpolarized and polarized proofs is that the former allows contractions at nearly any point in the proof while in the latter contraction only happens during some occurrences of the decide rules. As a result, we need to introduce some flexibility in how contexts are related between an unpolarized proof and the polarized proof emulating it. Given two multisets of LKF (LJF) formulas Γ and Γ' , we say that Γ' *extends* Γ if $FV(\Gamma) \subseteq FV(\Gamma')$ and every formula occurring in Γ also occurs in Γ' with the same or greater multiplicity – here, $FV(\Delta)$ denotes the set of variables occurring free in Δ . We say that an LKF (LJF) border sequent $\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'$ *extends an* LKF (LJF) *border sequent* $\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta$ if Γ' extends Γ and Δ' extends Δ .

Lemma 3. *Consider an application of an introduction rule in LK with concluding sequent S and premises S_1, \dots, S_p (for $p = 0, 1, 2$). Then for any LKF sequent S' that extends $[S]$, there exists a synthetic inference rule in LKF with conclusion S' and premises S'_1, \dots, S'_p such that for all $1 \leq i \leq p$, S'_i extends $[S_i]$. If we change LK to LJ and LKF to LJF above, this lemma also holds.*

PROOF. The proof proceeds by considering all the rules of LK . For example, consider the implication left-introduction rule for LK (see Figure 1)

$$\frac{\Gamma \vdash \Delta_1, A \quad B, \Gamma \vdash \Delta_2}{A \supset B, \Gamma \vdash \Delta_1, \Delta_2}.$$

Let the border sequent $\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'$ extend

$$[A \supset B]^l, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [\Delta_1]^r, [\Delta_2]^r.$$

Thus, Δ' extends both $[\Delta_1]^r$ and $[\Delta_2]^r$ and Γ' extends $[\Gamma]^l$ and contains the formula $[A \supset B]^l$. The following synthetic inference rule in LKF emulates this implication left-introduction rule.

$$\frac{\frac{\frac{\Gamma' \uparrow \cdot \vdash \cdot \uparrow [A]^r, \Delta'}{\Gamma' \uparrow \cdot \vdash [A]^r \uparrow \Delta'} S_r}{\Gamma' \uparrow \cdot \vdash \partial_-([A]^r) \uparrow \Delta'} \partial_-^r}{\Gamma' \vdash \partial_-([A]^r) \downarrow \Delta'} R_l}{\frac{\frac{\frac{\Gamma', [B]^l \uparrow \cdot \vdash \cdot \uparrow \Delta'}{\Gamma' \uparrow [B]^l \vdash \cdot \uparrow \Delta'} S_l}{\Gamma' \uparrow \partial_+([B]^l) \vdash \cdot \uparrow \Delta'} \partial_+^l}{\Gamma' \downarrow \partial_+([B]^l) \vdash \Delta'} R_l}{\Gamma' \downarrow \partial_-([A]^r) \supset \partial_+([B]^l) \vdash \Delta'} \supset_l}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} D_l}$$

Note that the border sequents $\Gamma' \uparrow \cdot \vdash \cdot \uparrow [A]^r, \Delta'$ and $[B]^l, \Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'$ extend $[\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [\Delta_1]^r, [A]^r$ and $[B]^l, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [\Delta_2]^r$, respectively. Observe that the stores in the leaves are possible since $[A]^r$ is positive (or atomic) and $[B]^l$ is negative (or atomic). A similar construction can be described for all of the other *LK* inference rules. Also, the proof that a similar lemma holds for intuitionistic logic proofs is proved in the same way.

Theorem 4. *Let Π be an *LK* derivation of a sequent S from the sequents S_1, \dots, S_n . Then there exists an *LKF* derivation Π' of $[S]$ from $[S_1], \dots, [S_n]$ such that each application in Π of a non-structural rule corresponds to a synthetic inference rule in Π' . If we replace *LK* with *LJ* and *LKF* with *LJF*, the resulting statement is also a theorem.*

PROOF. We proceed bottom-up by starting from the root of Π and build Π' by repeatedly applying Lemma 3. Since Lemma 3 is restricted to introduction rules, we need to consider here the contraction and initial rules in *LK*. The contraction rules in *LK* do not, in fact, translate to any rule or rules in *LKF*: the definition of “extends” makes this possible. The initial rule of *LK* is

$$\overline{P, \Gamma \vdash \Delta, P}$$

where P is atomic. The corresponding synthetic inference rules in *LKF* are either

$$\frac{\overline{\Gamma' \vdash P \Downarrow \Delta'}}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} \text{D}_r \quad \text{or} \quad \frac{\overline{\Gamma' \Downarrow P \vdash \Delta'}}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} \text{D}_l$$

depending on whether P has positive or negative polarity, respectively: also, Γ' and Δ' are extensions of $[\Gamma]^l$ and $[\Delta]^r$, respectively, containing $P = [P]^l = [P]^r$.

2.4.2. Mapping polarized proofs to unpolarized proofs

Given two multisets of *LKF* (*LJF*) formulas Γ and Γ' , we say that Γ' is a *contraction* of Γ if Γ and Γ' contain the same set of formulas but Γ can have more occurrences of them than in Γ' . We say that an *LKF* (*LJF*) border sequent $\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'$ is a *contraction of an *LKF* (*LJF*) border sequent* $\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta$ if Γ' is a contraction of Γ and Δ' is a contraction of Δ .

The following lemma is proved by analyzing the various synthetic inference rules that result when using decide-rules (D_l and D_r) on the result of using the left/right-translation on *LK* and *LJ* formulas.

Lemma 5. *Let S' be an *LKF* border sequent $\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'$ that is the left/right translation $[\cdot]$ of some *LK* sequent. Consider a synthetic inference rule in *LKF* with concluding sequent S' and premises S'_1, \dots, S'_p ($p \geq 0$). It is the case that $0 \leq p \leq 2$ and that there exists*

1. an *LK* sequent S , such that S' is a contraction of $[S]$, and
2. an *LK* rule application with conclusion S and premises S_1, \dots, S_p such that for all $0 \leq i \leq p$, S'_i is a contraction of $[S_i]$.

If we change LK to LJ and LKF to LJF above, this lemma remains true.

Theorem 6. *Let Π' be a proof of a border sequent S' in LKF such that $S' = [S]$ for some LK -sequent S . Then there exists an LK -proof Π of S such that each synthetic inference rule in Π' corresponds to a single rule application in Π . If we change LK to LJ and LKF to LJF above, this lemma remains true.*

PROOF. We proceed top-down starting from the leaves of Π' and build Π by repeatedly applying Lemma 5. At each step, we get as the conclusion of an LK rule application a sequent S^* such that the one obtained in the corresponding step of Π' is a contraction of $[S^*]$.

As described in [22], comparing two proof systems can be done at three different levels of *adequacy*. *Relative completeness* claims that the provable formulas are the same in the two proof systems. *Full completeness of proofs* claims that complete proofs of theorems are in one-to-one correspondence between the two proof systems. Finally, the most demanding notion of adequacy is *full completeness of derivations* which claims that (open) derivations (such as inference rules themselves) are also in one-to-one correspondence between the two proof systems. What Theorems 4 and 6 imply is that we have this strongest form of adequacy on derivations, where one step in LK or in LJ corresponds to one synthetic inference rule in LKF or in LJF , respectively.

3. Synthetic inference rules for bipolar theories

If we limit the alternation of polarity within a negative formula to just one flip, then the synthetic inference rules for such a formula are particularly simple. In fact, such synthetic rules do not explicitly mention logical connectives.

Definition 7 (Bipole for B). *Let B be a polarized negative formula in either LKF or LJF . A bipole for B is a synthetic inference rule for B (see Definition 2) in which all formulas stored using the store rules (S_l, S_r among the inference rules justifying this synthetic inference rule) are atomic formulas.*

Bipoles are, therefore, synthetic inference rules in which the only difference between the concluding border sequent and any one of its premises is the presence or absence of atomic formulas.

Example 8. *Let $P_1(x)$, $P_2(x)$, $Q(x)$, and $R(x, y)$ be positive atomic formulas and assume that the polarized formula $\forall x(((P_1(x) \supset P_2(x)) \wedge^+ Q(x)) \supset \exists y R(x, y))$ is a member of Γ' . The following LKF derivation justifies a synthetic inference*

rule for this formula.

$$\begin{array}{c}
\frac{\Gamma', P_1(t) \uparrow \cdot \vdash \cdot \uparrow P_2(t), \Delta'}{\Gamma', P_1(t) \uparrow \cdot \vdash \cdot \uparrow P_2(t) \uparrow \Delta'} S_r \\
\frac{\Gamma', P_1(t) \uparrow \cdot \vdash \cdot \uparrow P_2(t) \uparrow \Delta'}{\Gamma' \uparrow P_1(t) \vdash P_2(t) \uparrow \Delta'} S_l \\
\frac{\Gamma' \uparrow P_1(t) \vdash P_2(t) \uparrow \Delta'}{\Gamma' \uparrow \cdot \vdash P_1(t) \supset P_2(t) \uparrow \Delta'} \supset_r \\
\frac{\Gamma' \uparrow \cdot \vdash P_1(t) \supset P_2(t) \uparrow \Delta'}{\Gamma' \vdash P_1(t) \supset P_2(t) \downarrow \Delta'} R_r \\
\frac{\Gamma' \vdash P_1(t) \supset P_2(t) \downarrow \Delta'}{\Gamma' \vdash (P_1(t) \supset P_2(t)) \wedge^+ Q(t) \downarrow \Delta'} \wedge_r^+ \\
\frac{\Gamma' \vdash (P_1(t) \supset P_2(t)) \wedge^+ Q(t) \downarrow \Delta'}{\Gamma' \downarrow \forall x((P_1(x) \supset P_2(x)) \wedge^+ Q(x)) \supset \exists y R(x, y) \uparrow \Delta'} \forall_l \\
\frac{\Gamma' \downarrow \forall x((P_1(x) \supset P_2(x)) \wedge^+ Q(x)) \supset \exists y R(x, y) \uparrow \Delta'}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} D_l \\
\frac{\Gamma', R(t, z) \uparrow \cdot \vdash \cdot \uparrow \Delta'}{\Gamma' \uparrow R(t, z) \vdash \cdot \uparrow \Delta'} S_l \\
\frac{\Gamma' \uparrow R(t, z) \vdash \cdot \uparrow \Delta'}{\Gamma' \uparrow \exists y R(t, y) \vdash \cdot \uparrow \Delta'} \exists_l \\
\frac{\Gamma' \uparrow \exists y R(t, y) \vdash \cdot \uparrow \Delta'}{\Gamma' \downarrow \exists y R(t, y) \vdash \Delta'} R_l \\
\frac{\Gamma' \downarrow \exists y R(t, y) \vdash \Delta'}{\Gamma' \downarrow \exists y R(t, y) \vdash \Delta'} \supset_l
\end{array}$$

Note that since we are assuming that atomic polarity assignments are stable under substitution, it must be the case that the atoms $P_1(t)$, $P_2(t)$, $Q(t)$, $R(t, z)$ are all positive.¹ In order to apply the rule I_r in this derivation, it must be the case that $Q(t) \in \Gamma'$. Thus, the corresponding bipole in LKF is

$$\frac{P_1(t), Q(t), \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta, P_2(t) \quad R(t, z), Q(t), \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta}{Q(t), \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta}$$

If we were to build the same kind of synthetic inference rule in LJF, the corresponding bipole would be

$$\frac{P_1(t), Q(t), \Gamma \uparrow \cdot \vdash \cdot \uparrow P_2(t) \quad R(t, z), Q(t), \Gamma \uparrow \cdot \vdash \cdot \uparrow E}{Q(t), \Gamma \uparrow \cdot \vdash \cdot \uparrow E} .$$

In both rules, the variable z does not occur in the conclusion.

Following the classification of formulas in intuitionistic linear logic given in [7], we organize polarized first-order classical and intuitionistic formulas into a hierarchy based on the alternation of polarized connectives.

Definition 9 (Hierarchy of polarized formulas). We define the following hierarchy of negative and positive classical formulas (denoted \mathcal{N}^C and \mathcal{P}^C , respectively). The classes \mathcal{N}_0^C and \mathcal{P}_0^C are both equal to the set of atomic formulas. The rest of the hierarchy is defined recursively as follows:

$$\begin{aligned}
\mathcal{N}_{n+1}^C &::= \forall x \mathcal{N}_{n+1}^C \mid \mathcal{N}_{n+1}^C \wedge^- \mathcal{N}_{n+1}^C \mid \mathcal{N}_{n+1}^C \vee^- \mathcal{N}_{n+1}^C \mid \mathcal{P}_{n+1}^C \supset \mathcal{N}_{n+1}^C \mid \mathcal{P}_n^C \mid t^- \mid f^- \\
\mathcal{P}_{n+1}^C &::= \exists x \mathcal{P}_{n+1}^C \mid \mathcal{P}_{n+1}^C \wedge^+ \mathcal{P}_{n+1}^C \mid \mathcal{P}_{n+1}^C \vee^+ \mathcal{P}_{n+1}^C \mid \mathcal{N}_n^C \mid t^+ \mid f^+
\end{aligned}$$

The hierarchy of negative and positive intuitionistic formulas (denoted \mathcal{N}^I , \mathcal{P}^I , respectively) is defined analogously, by simply omitting the cases of \vee^- and f^- in the definition of \mathcal{N}_{n+1}^I . Also, in LJF, the classes \mathcal{N}_0^I and \mathcal{P}_0^I are both equal to the set of atomic formulas.

¹Sonia: P_2 does need to be negative to be stored on the right?

Definition 10 (Bipolar formula). Any formula in the class \mathcal{N}_2^C is a classical bipolar formula. Any formula in the class \mathcal{N}_2^I is an intuitionistic bipolar formula.

Example 11. The formula $\forall x(((P_1(x) \supset P_2(x)) \wedge^+ Q(x)) \supset \exists yR(x, y))$ under focus in the conclusion of the derivation of Example 8 can be read as both a classical and an intuitionistic bipolar formula.

We make the following useful additional definitions. A formula occurrence in a sequent in *LKF* or in *LJF* is *in the inner zone* if that occurrence appears between either \uparrow or \downarrow and \vdash . An *LKF* polarized formula B is *level- n* (for $n \geq 0$) if it is a member of $\mathcal{N}_n^C \cup \mathcal{P}_n^C$. An *LKF*-sequent is *level- n* if every formula in its inner zone is of level- n . (Note that a border sequent is of level- n for all natural number n .) The *size* of an *LKF*-sequent is the total number of occurrences of logical connectives in formulas appearing in that sequent's inner zone. Note that the definitions for level and size can be extended directly to the intuitionistic case by defining level- n using $\mathcal{N}_n^I \cup \mathcal{P}_n^I$ and by considering *LJF* sequents instead of *LKF* sequents.

Theorem 12. A synthetic inference rule for a bipolar formula is a bipole.

PROOF. We restrict our attention first to the *LJF* proof system. Let B be a polarized negative formula of level-2 and consider a synthetic inference rule for $B \in \Gamma$ of the sequent $\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta$. The last inference rule of this synthetic inference rule is D_l with premise $\Gamma \downarrow B \vdash \Delta$. As we move up within the synthetic inference rule from this sequent, we first move through synchronous introduction rules until we arrive at a release rule. At that point, the single formula in that sequent is of level 1. During the asynchronous phase, any instances of store rules will store formulas of level 0: in other words, the only formulas stored in this synthetic inference rule are atomic formulas. Hence, this synthetic inference rule is a bipole. The same argument can be made for classical bipolar formulas and synthetic inference rules in *LKF*.

The following is the converse of Theorem 12.

Theorem 13. If every synthetic inference rule for a given negative formula is a bipole then that formula is bipolar.

PROOF. We first consider *LJF* proofs. We say that a formula has minimal level k if it has level k but does not have level $k - 1$ (for the base case, atoms have minimal level 0). Assume that every synthetic inference rule of the negative polarized formula B is a bipole but that B is not a bipolar formula. Hence there exists $k \geq 3$ such that B is of minimal level k . Consider a synthetic inference rule for $B \in \Gamma$ of the sequent $\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta$. During the construction of the synchronous phase (from conclusion to premise), it is possible to always pick instances of the synchronous introduction rules so that whenever the inner formula is positive, its minimal level is k and if it is negative, its minimal level is $k - 1$. During the construction of the asynchronous phase, among the formulas stored there

must be one that is negative and has minimal order $k - 2$. Since $k \geq 2$ that stored formula is not atomic, and, hence, this synthetic inference rule is not a bipole, which is a contradiction. This proof can easily be extended to handle *LKF* proofs.

Which synthetic rules correspond to a given bipolar formula B can be computed directly from that formula using the *LKF* and *LJF* proof systems. In general, such synthetic rules have the form

$$\frac{\overline{\Gamma}_1, \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta, \overline{\Delta}_1 \quad \cdots \quad \overline{\Gamma}_r, \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta, \overline{\Delta}_r}{\overline{\Gamma}_0, \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta, \overline{\Delta}_0} .$$

Here, $r \geq 0$ and

1. the schematic variables for this rule are Γ and Δ and these range over multisets of formulas,
2. for $0 \leq i \leq r$, the multisets $\overline{\Gamma}_i, \overline{\Delta}_i$ are atomic subformulas of B ,
3. all free variables in these subformulas range over first-order terms, and
4. all the eigenvariables in $\overline{\Gamma}_j, \overline{\Delta}_j$ introduced in Π by \forall_r or \exists_l do not occur free in the conclusion.

The intuitionistic version of the rule above is required to satisfy the usual restriction that at most one formula is allowed on the right side of a sequent. Note that synthetic inference rules do not explicitly mention logical connectives, only atomic formulas.

In general, the computation of a synthetic inference rule from a bipolar formula starts by trying to build a bipole in which the (left) focus is on the formula B . As we make choices in which rules to apply during the synchronous phase, we might need to deal with a focus on an atomic formula occurrence, say A . If A is negatively biased and focused on the left, then A must also be present on the right-hand-side, which is possible if $A \in \overline{\Delta}_0$; dually, if A is positively biased and focused on the right, then A must also be present on the left-hand-side, which is possible if $A \in \overline{\Gamma}_0$. If, however, A is negatively biased and focused on the right or is positively biased and focused on the left, a release rule must be used and we transition to the asynchronous phase. As we continue constructing the asynchronous phase, all atoms appearing in the inner zone will be stored on either the left or right and, hence, they will populate either Γ_i or Δ_i depending on which branch that store occurs.

This iterative process for computing bipoles from bipolar formulas is easily seen to be terminating using the following measure. We assign to every *LJF* and *LKF* sequent S a triple $\langle m, n, p \rangle$ of natural numbers as follows: m is the total number of logical connectives occurring in formulas in the inner zone; n is the number of formulas in the inner zone; and p is 0 if the sequent is an \uparrow sequent and 1 if the sequent is a \downarrow sequent. We shall refer to this triple as the *measure* of a sequent and we use the usual lexicographic ordering on triples to provide a well-ordering on this measure. Notice that when moving from the conclusion to a premise for every rule except the decide rules (D_l, D_r), the measure of the

sequent gets smaller. In particular, the first component gets smaller for any introduction rule, the second component gets smaller for any store rule (S_l , S_r), and the third component gets smaller for the release rules (R_l , R_r).

Appendix A contains a λ Prolog [23, 24] executable specification of a predicate that relates a bipolar formula to its various bipoles. Given the nature of λ Prolog, this specification is both compact and explicit about the scope of bindings for schematic variables and eigenvariables. The termination of this λ Prolog specification is easily shown using the measure of sequents mentioned above.

Our main project in this paper is to have a general method for extending both LK and LJ with inference rules that capture certain classes of axioms. From what we have seen in this section, we can now make three observations concerning this project. The first is that by restricting to bipolar formulas, these new inference rules involve only atomic formulas (and schematic variables ranging over contexts). The second is that working with axioms as unpolarized formulas is not sufficient: in order to construct inference rules, we need to start with *polarized* axioms. Finally, a polarized axiom B that is a positive bipolar formula is logically equivalent to the negative bipolar $\partial_-(B)$. Hence we may consider, without loss of generality, that axioms in any theory \mathcal{T} are negative formulas.

Definition 14 (Rules from polarized axioms). *Let $\langle \delta, \mathcal{T} \rangle$ be a finite set of bipolar formulas. We define $LK\langle \delta, \mathcal{T} \rangle$ to be the extension of LK with inference rules derived from the polarized theory $\langle \delta, \mathcal{T} \rangle$ as follows. For every $B \in \mathcal{T}$ and every synthetic inference rule for B , say,*

$$\frac{\Gamma_1 \uparrow \cdot \vdash \cdot \uparrow \Delta_1 \quad \dots \quad \Gamma_n \uparrow \cdot \vdash \cdot \uparrow \Delta_n}{\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta},$$

we place in $LK\langle \delta, \mathcal{T} \rangle$ the inference rule

$$\frac{\Gamma_1 \vdash \Delta_1 \quad \dots \quad \Gamma_n \vdash \Delta_n}{\Gamma \vdash \Delta}.$$

Since $\Gamma, \Gamma_1, \dots, \Gamma_n, \Delta, \Delta_1, \dots, \Delta_n$ are composed of either atomic formulas or schematic (context) variables, there are no logical connectives that need to be “de-polarize” when moving from the first inference rule above to the second.

Example 15. *Let δ be any atom bias assignment that assigns positive to the open atomic formulas $P_1(x), P_2(x), Q(x), R(x, y)$ and all their instances (see Example 8). If the polarized formula*

$$\forall x(((P_1(x) \supset P_2(x)) \wedge^+ Q(x)) \supset \exists y R(x, y))$$

is a member of \mathcal{T} , then the following rule is contained in $LK\langle \delta, \mathcal{T} \rangle$

$$\frac{P_1(t), Q(t), \Gamma \vdash \Delta, P_2(t) \quad R(t, z), Q(t), \Gamma \vdash \Delta}{Q(t), \Gamma \vdash \Delta}$$

while the following rule is contained in $LJ\langle\delta, \mathcal{T}\rangle$.

$$\frac{P_1(t), Q(t), \Gamma \vdash P_2(t) \quad R(t, z), Q(t), \Gamma \vdash E}{Q(t), \Gamma \vdash E}$$

In both of these rules, the eigenvariable z does not occur in the conclusion of the corresponding rule.

We shall now prove that cut is an admissible rule in the $LK\langle\delta, \mathcal{T}\rangle$ and $LJ\langle\delta, \mathcal{T}\rangle$ proof systems, where the cut frees for these systems are, respectively,

$$\frac{\Gamma \vdash \Delta, A \quad A, \Gamma \vdash \Delta}{\Gamma \vdash \Delta} \text{ cut}_{LK} \quad \text{and} \quad \frac{\Gamma \vdash A \quad A, \Gamma \vdash B}{\Gamma \vdash B} \text{ cut}_{LJ}.$$

Our proof here will be simple since we can directly use the cut-elimination theorem for LJF and LKF given in [3, 21].

Theorem 16 (Cut admissibility for $LJ\langle\delta, \mathcal{T}\rangle$). *Let $\langle\delta, \mathcal{T}\rangle$ be a set of bipolar formulas. The cut rule is admissible for the proof systems $LJ\langle\delta, \mathcal{T}\rangle$.*

PROOF. The following two cut rules (among others) are proved to be admissible for LJF in [3].

$$\frac{\Gamma_1 \uparrow \Theta_1 \vdash \cdot \uparrow P \quad \Gamma_2 \uparrow P, \Theta_2 \vdash \mathcal{R}}{\Gamma_1, \Gamma_2 \uparrow \Theta_1, \Theta_2 \vdash \mathcal{R}} \text{ Cut}^+ \quad \frac{\Gamma_1 \uparrow \Theta_1 \vdash C \uparrow \cdot \quad \Gamma_2, C \uparrow \Theta_2 \vdash \mathcal{R}}{\Gamma_1, \Gamma_2 \uparrow \Theta_1, \Theta_2 \vdash \mathcal{R}} \text{ Cut}^-$$

Here, P is a positive formula, C is a negative formula or a positive atom, and (as in Figure 3) \mathcal{R} denotes $\Delta_1 \uparrow \Delta_2$ where the union of Δ_1 and Δ_2 contains at most one formula. We shall apply these admissibility results for LJF to immediately yield the cut-admissibility result for $LJ\langle\delta, \mathcal{T}\rangle$.

Assume that we have (cut-free) proofs in $LJ\langle\delta, \mathcal{T}\rangle$ of the sequents $\Gamma \vdash B$ and $B, \Gamma \vdash E$. Using Theorem 4 and Definition 14, we have proofs in LJF of

$$\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [B]^r \quad \text{and} \quad \mathcal{T}, [B]^l, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [E]^r.$$

Consider the following two cases.

Case 1: B is atomic. In this case, $[B]^r$ and $[B]^l$ are equal to B . By using the admissibility of Cut^- , we know that $\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [E]^r$. We cannot apply Cut^- directly, but it is easy to see that for atomic formula B (of either polarity), $\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow B$ is provable in LJF if and only if $\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash B \uparrow \cdot$ is provable. We can now apply the admissibility of Cut^- to derive the sequent $\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [E]^r$.

Case 2: B is not atomic. In this case, $[B]^r$ and $[B]^l$ are different: the first has positive polarity while the second has negative polarity. Since these formulas are different, a focused cut rule would not apply directly to them. However, using the completeness theorem for LJF mentioned in Section 2.3 and proved in [3], since $B \supset B$ is provable in LJ then every polarization of $B \supset B$ is provable in LJF . Thus, it must be the case that we have an LJF proof of $\cdot \uparrow \vdash [B]^r \supset [B]^l \uparrow \cdot$ and (by inversion) $\cdot \uparrow [B]^r \vdash [B]^l \uparrow \cdot$. By the admissibility of Cut^- , we can

conclude that there is a (cut-free) *LJF* proof of $\mathcal{T}, [\Gamma]^l \uparrow [B]^l \vdash \cdot \uparrow [E]^r$. By the admissibility of *Cut*⁺, we can conclude that there is a (cut-free) *LJF* proof of $\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [E]^r$.

Thus, in either case, we have $\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [E]^r$. By Theorem 6 and Definition 14, we can conclude that $\Gamma \vdash E$ has a cut-free *LJ*(δ, \mathcal{T}).

Theorem 17 (Cut admissibility for *LK*(δ, \mathcal{T})). *Let $\langle \delta, \mathcal{T} \rangle$ be a polarized, geometric theory. The cut rule is admissible for the proof systems *LK*(δ, \mathcal{T}).*

PROOF. Consider the following two cut rules

$$\frac{\frac{\Gamma_1 \uparrow \Theta_1 \vdash \Omega_1 \uparrow P, \Delta_1 \quad \Gamma_2 \uparrow P, \Theta_2 \vdash \Omega_2 \uparrow \Delta_2}{\Gamma_1, \Gamma_2 \uparrow \Theta_1, \Theta_2 \vdash \Omega_1, \Omega_2 \uparrow \Delta_1, \Delta_2}}{\Gamma_1 \uparrow \Theta_1 \vdash C, \Omega_1 \uparrow \Delta_1 \quad \Gamma_2, C \uparrow \Theta_2 \vdash \Omega_2 \uparrow \Delta_2} \quad \frac{\Gamma_1, \Gamma_2 \uparrow \Theta_1, \Theta_2 \vdash \Omega_1, \Omega_2 \uparrow \Delta_1, \Delta_2}{\Gamma_1, \Gamma_2 \uparrow \Theta_1, \Theta_2 \vdash \Omega_1, \Omega_2 \uparrow \Delta_1, \Delta_2}}$$

Here, P is a positive formula, C is a negative formula or a positive atom. The admissibility of these rules in *LKF* follows from the cut-admissibility result in [21] (these two cuts correspond to the $dcut_f$ rule in the one-sided variant of *LKF* used in that paper). We shall apply the cut-admissibility result for *LKF* to immediately yield the cut-admissibility result for *LK*(δ, \mathcal{T}).

Assume that we have (cut-free) proofs in *LK*(δ, \mathcal{T}) of the sequents $\Gamma \vdash B, \Delta$ and $B, \Gamma \vdash \Delta$. Using Theorem 4 and Definition 14, we have proofs in *LKF* of

$$\mathcal{T}, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [B]^r, [\Delta]^r \quad \text{and} \quad \mathcal{T}, [B]^l, [\Gamma]^l \uparrow \cdot \vdash \cdot \uparrow [\Delta]^r.$$

The rest of this proof follows the same steps as the proof regarding *LJ*(δ, \mathcal{T}) (Theorem 16).

Let B be a first-order formula. If $\langle \delta, \hat{B} \rangle$ is an *LJF*-polarization of B then it is also an *LKF*-polarization of B since there are more options for polarizing classical formulas. The converse obviously does not hold. Moreover, the set of formulas which can be polarized as a bipolar is strictly greater in the classical setting. Indeed, for any atomic bias assignment, the formula $(P_1 \supset P_2) \vee (Q_1 \supset Q_2)$ can be polarized classically to yield the bipolar $(P_1 \supset P_2) \vee (Q_1 \supset Q_2)$ while there is no polarization of this formula that gives rise to an intuitionistic bipolar formula.

4. Synthetic inference rules for geometric theories

The quest of generating sequent proof calculi for a large class of axiomatic extensions of classical and intuitionistic logics has been focus of attention for quite some time now. As it is well-known, simply adding an axiom as a theorem in the logical system is not a solution, since the resulting system may not yield the same theorems of the extended logic [25, 26].

Such a problem can be overcome by converting axioms of a certain shape into rules of sequent calculus in such a way that the logical content of the axiom is replaced by the meta-linguistic meaning of sequent rules [5, 9, 12, 10, 11, 27]. In this section, we will show how bipolar formulas and focusing provide a generalization of such a method.

4.1. Geometric axioms as bipolar formulas

There are many examples of geometric theories in different areas of logic and mathematics, such as topology and category theory [6]. Since geometric axioms form a proper subclass of bipolars, the approach developed in this paper can be applied for translating this class of axioms into synthetic inference rules.

Definition 18. A geometric implication is a first-order formula having the form

$$\forall \bar{z}(P_1 \wedge \dots \wedge P_m \supset \exists \bar{x}_1 M_1 \vee \dots \vee \exists \bar{x}_n M_n),$$

where each P_i is an atomic formula, each M_j is a conjunction of atomic formulas $Q_{j_1}, \dots, Q_{j_{k_j}}$, and none of the variables in the lists $\bar{x}_1, \dots, \bar{x}_n$ are free in P_i . A geometric theory is a finite set of geometric implications. We shall also assume that if the list of variables \bar{x}_i is empty then M_i is just an atom: otherwise, this formula can be written as a conjunction of geometric implications. For example, if \bar{x}_1 is empty, then the following equivalence holds

$$\forall \bar{z}(\mathbf{P} \supset (\bigwedge_{l=1}^{j_1} Q_{1l}) \vee \mathbf{M}) \equiv \bigwedge_{l=1}^{j_1} (\forall \bar{z}(\mathbf{P} \supset (Q_{1l} \vee \mathbf{M})))$$

A coherent implication [6] is the universal closure of implications of the form $D_1 \supset D_2$, where D_i is built up from atoms using conjunction, disjunction and existential quantification. It is routine to check that coherent implications are intuitionistically equivalent to conjunctions of geometric implications (see e.g., [6, Proposition 2.6]).

As one can expect, different polarizations of geometric implications can give rise to different bipoles. Note that if we wish to polarize the displayed formula in Definition 18, the conjunctions within M_i must be polarized positively (assuming that the list of variable \bar{x}_i is a non-empty). Hence, the polarized geometric implication

$$\forall \bar{z}(P_1^\pm \wedge^\pm \dots \wedge^\pm P_n^\pm \supset \exists \bar{x}_1 \hat{M}_1 \vee^\pm \dots \vee^\pm \exists \bar{x}_n \hat{M}_n)$$

is a bipolar formula, where X^\pm means that X may have any polarity and $\hat{M}_j = Q_{j_1}^\pm \wedge^+ \dots \wedge^+ Q_{j_{k_j}}^\pm$.

As an example, consider the polarization of formulas such that \wedge and \vee are replaced by their positive versions, the atoms P_i ($1 \leq i \leq n$) are assigned a positive bias, and all the atoms Q_{j_l} are given any polarization (for all and $1 \leq j \leq n, 1 \leq j_l \leq k_j$). A synthetic rule corresponding to a bipole for this formula is

$$\frac{\overline{Q_1}[\bar{y}_1/\bar{x}_1], \overline{P}, \Gamma \vdash \Delta \quad \dots \quad \overline{Q_n}[\bar{y}_n/\bar{x}_n], \overline{P}, \Gamma \vdash \Delta}{\overline{P}, \Gamma \vdash \Delta} \text{GRS}$$

which is justified by a derivation with the following structure:

$$\frac{\frac{\frac{\overline{Q}_1 \rho_1, \overline{P}, \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta}{\overline{P}, \Gamma \uparrow M_1 \rho_1 \vdash \cdot \uparrow \Delta} \wedge_l^+, S_l \quad \frac{\overline{Q}_n \rho_n, \overline{P}, \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta}{\overline{P}, \Gamma \downarrow M_n \rho_n \vdash \Delta} \wedge_l^+, S_l}{\overline{P}, \Gamma \vdash \bigwedge^+ P_i \downarrow \Delta} \wedge_r^+, I_r \quad \frac{\overline{P}, \Gamma \downarrow M_n \rho_n \vdash \Delta}{\overline{P}, \Gamma \downarrow \bigvee^+ \exists \overline{x}_j M_j \vdash \Delta} R_r, \vee_l^+, \exists_l}{\overline{P}, \Gamma \downarrow \forall z. (\bigwedge^+ P_i \supset \bigvee^+ \exists \overline{x}_j M_j) \vdash \Delta} \forall_l, \supset_l$$

where the variable renaming substitution ρ_i is equal to $[\overline{y}_i/\overline{x}_i]$, the symbols \overline{Q}_j and \overline{P} denote the multisets of atomic formulas $Q_{j_1}, \dots, Q_{j_{k_j}}$ and P_1, \dots, P_m , respectively, and the eigenvariables in the lists $\overline{y}_1, \dots, \overline{y}_n$ do not occur free in the conclusion. In [10, 27, 28], the *GRS* synthetic inference rule is called the *geometric rule scheme*.

Another class of formulas described in [27, Chapter 5] are the *co-geometric axioms*, which are of the form

$$\forall z (\forall \overline{x}_1 M_1 \wedge \dots \wedge \forall \overline{x}_n M_n \supset P_1 \vee \dots \vee P_m),$$

where M_j is the disjunction of atoms $Q_{j_1} \vee \dots \vee Q_{j_{k_j}}$ (for $1 \leq j \leq n$). If the disjunctions on the right of the implication are polarized negatively, then the polarized axiom can be a bipolar formula. In particular, the polarized formula

$$\forall z (\forall \overline{x}_1 M_1 \wedge^\pm \dots \wedge^\pm \forall \overline{x}_n M_n \supset P_1 \vee \dots \vee P_m),$$

with $M_j = Q_{j_1} \vee \dots \vee Q_{j_{k_j}}$ and P_i polarized negatively, gives rise to the synthetic inference rule in *LKF* (called *co-geometric rule scheme* in [27])

$$\frac{\Gamma \vdash \overline{Q}_1[\overline{y}_1/\overline{x}_1], \overline{P}, \Delta \quad \dots \quad \Gamma \vdash \overline{Q}_n[\overline{y}_n/\overline{x}_n], \overline{P}, \Delta}{\Gamma \vdash \overline{P}, \Delta} \text{co-GRS}_c$$

If we restrict ourselves to the intuitionistic case, we must have $m = 1$ and $M_j = Q_j$ for $1 \leq j \leq n$. The synthetic rule in this case is given by

$$\frac{\Gamma \vdash Q_1[\overline{y}_1/\overline{x}_1] \quad \dots \quad \Gamma \vdash Q_n[\overline{y}_n/\overline{x}_n]}{\Gamma \vdash P} \text{co-GRS}_i.$$

Given a (classical or intuitionistic) geometric/co-geometric theory T , a complete proof calculus for it can be obtained by adding to an appropriate base (classical or intuitionistic) proof system the rules that follow the scheme corresponding to the (polarized) axioms in T . Observe that our setting avoids the *closure condition rules* in [9] since contraction is implicit in the focused setting. Moreover, all the structural properties of the basic sequent calculi are preserved by the addition of rules following the schemes given above.

As the following example illustrates, not all bipolar formulas are geometric or co-geometric formulas.

Example 19. *In set theory, the following implication relates the subset and membership predicates:*

$$\forall yz. (\forall x (x \in y \supset x \in z) \supset y \subseteq z).$$

This formula yields a bipolar formula in both *LKF* and *LJF* under any polarization of the binary atomic predicates \in and \subseteq . Assuming that these predicates are given positive polarity, the corresponding *LJF*-synthetic inference rule is

$$\frac{x \in y, \Gamma \vdash x \in z \quad y \subseteq z, \Gamma \vdash E}{\Gamma \vdash E} .$$

Assuming that these predicates are given negative polarity, the corresponding *LJF*-synthetic inference rule is

$$\frac{x \in y, \Gamma \vdash x \in z}{\Gamma \vdash y \subseteq z} .$$

In both of these synthetic inference rules, x is an eigenvariable for that rule.

4.2. Universal axioms as bipoles

A *universal implication* is a restricted geometric formula of the form

$$\forall z (P_1 \wedge \dots \wedge P_m \supset Q_1 \vee \dots \vee Q_n)$$

where all the P_i and Q_i are atomic formulas. This subset of geometric implications allows for more choices in the selection of polarities while still remaining bipolar formulas. In fact, polarized universal implications (in *LKF*) can have the form

$$\forall z (P_1^\pm \wedge^\pm \dots \wedge^\pm P_m^\pm \supset Q_1^\pm \vee^\pm \dots \vee^\pm Q_n^\pm).$$

Consider the *LKF*-polarized version of the universal implication with the connectives \wedge and \vee and the atoms Q_1, \dots, Q_n all polarized negatively. The correspondent synthetic rule is

$$\frac{\Gamma \vdash P_1, \bar{Q}, \Delta \quad \dots \quad \Gamma \vdash P_m, \bar{Q}, \Delta}{\Gamma \vdash \bar{Q}, \Delta} RR_c$$

is called the *right universal rule scheme* in [28]. In the intuitionistic setting, we can use this polarization only when $n = 1$, in which case the *LJF* synthetic inference rule is

$$\frac{\Gamma \vdash P_1 \quad \dots \quad \Gamma \vdash P_m}{\Gamma \vdash Q} RR_i.$$

Figure 4 presents four synthetic rules in the classical setting, corresponding to different possible polarity assignments. For example, RL_c is given by the following polarizations: $\{\wedge^+, \vee^+, P_i^+, Q_j^\pm\}$ and $\{\wedge^+, \vee^-, P_i^+, Q_j^+\}$.

A special case of universal implications are *Horn clauses*, that is, formulas of the form $\forall z (P_1 \wedge \dots \wedge P_m \supset Q)$, where where all the P_i and Q are atomic formulas. As such, a Horn clause is also a universal implication and the various polarizations described above can be applied to them, yielding different synthetic inference rules. In particular, if we polarize atoms and conjunctions negatively

$$\begin{array}{c}
\frac{\Gamma \vdash P_1, \overline{Q}, \Delta \quad \dots \quad \Gamma \vdash P_m, \overline{Q}, \Delta}{\Gamma \vdash \overline{Q}, \Delta} \quad RR_c \qquad \frac{Q_1, \overline{P}, \Gamma \vdash \Delta \quad \dots \quad Q_n, \overline{P}, \Gamma \vdash \Delta}{\overline{P}, \Gamma \vdash \Delta} \quad RL_c \\
\frac{\Gamma \vdash P_1, \Delta \quad \dots \quad \Gamma \vdash P_m, \Delta \quad Q_1, \Gamma \vdash \Delta \quad \dots \quad Q_n, \Gamma \vdash \Delta}{\Gamma \vdash \Delta} \quad SS_c \qquad \frac{}{\overline{P}, \Gamma \vdash \overline{Q}, \Delta} \quad BB_c
\end{array}$$

Figure 4: Synthetic (classical) rule schemes corresponding to different polarizations of universal implications.

in the Horn clause above, the resulting *LJF*-synthetic inference rule is the back-chaining rule

$$\frac{\Gamma \vdash P_1 \quad \dots \quad \Gamma \vdash P_m}{\Gamma \vdash Q} \quad BC.$$

On the other hand, if we polarize atoms and conjunctions positively in the Horn clause above, the resulting *LJF*-synthetic inference rule is the forward-chaining rule

$$\frac{\Gamma, P_1, \dots, P_m, Q \vdash B}{\Gamma, P_1, \dots, P_m \vdash B} \quad FC.$$

Finally, it is worth noting that, in the classical setting, it is possible to extend the first-order language with new function symbols, so that any axiom can be converted to a finite set of coherent implications [6]. We illustrate such a method next.

Example 20. *Given a first order binary relation R , a $\langle i, j, m, n \rangle$ -convergency axiom (see [12]) has the form*

$$\forall xyz((R^i(x, y) \wedge R^j(x, z)) \supset \exists u.(R^m(y, u) \wedge R^n(z, u)))$$

where $R^0(x, y)$ is defined to be $x = y$ and $R^{i+1}(x, y)$ is defined to be $\exists v.R(x, v) \wedge R^i(v, y)$ for $i \geq 1$. As noted before, such formulas are bipolars iff conjunctions in the head of the implication are polarized positively. This restriction can be bypassed using skolemization. In fact, by prenexing quantifiers and then skolemizing the remaining existential quantifiers, convergency axioms are transformed into a set of Horn (relational) formulas of the form

$$\forall \bar{z}(R(s_1, t_1) \wedge \dots \wedge R(s_m, t_m) \supset R(s_0, t_0))$$

where s_i, t_i are terms built from \bar{z} and Skolem function constants.

Convergency axioms generalize Scott-Lemmon axioms [29] (a.k.a. Geach axioms), which correspond to a “confluence” condition on the relational structure of modal logic (see Section 5).

In the next section we will shed some light on the behavior of axioms falling outside the boundary of bipolar formulas.

4.3. Beyond bipoles

Theorems 12 and 13 set a boundary to the process of transforming axioms into rules in the classical/intuitionistic settings, since they identify *the exact class* of formulas that can be seen as synthetic inference rules.² In this section, we illustrate how to relate non-bipolars to the *systems of rules* formalism, introduced in [28]. Such formalism is an extension of the *axioms-as-rules formalism*, since it allows for different sequent rules connected by conditions on the order of their applicability and with the possibly of sharing meta-variables for formulas or sets of formulas. While in [28] systems of rules were applied to the class of generalized geometric implications, in [30] a connection between hypersequents and a subclass of systems of rules is shown for *propositional* intermediate logics.

We will only illustrate how the systems of rules method would work for some chosen examples, since a complete discussion of the subject would fall out the scope of the present paper.

Example 21. *The powerset axiom in set theory, written as*

$$A = \forall z \exists w \forall y (\forall x ((x \in y) \supset (x \in z)) \supset (y \in w)),$$

is not bipolar due to the alternation of the positive and negative quantifiers. It is also not a generalized geometric implication, since it has a negative occurrence of implication. If we write $B(z, w)$ for $\forall y (\forall x ((x \in y) \supset (x \in z)) \supset (y \in w))$, a focus on A would justify the following inference rule

$$\frac{B(s, w), \Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta}{\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta} \text{ decide on } A, \forall_l, R_l, \exists_l$$

where s is the substitution instance of $\forall z$ and w is not free in s nor in Γ, Δ . Now, $B(s, w)$ is bipolar with corresponding bipole

$$\frac{x \in y, \Gamma' \uparrow \cdot \vdash \cdot \uparrow x \in s, \Delta' \quad y \in w, \Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} \text{ decide on } B(s, w), \text{ etc.}$$

Here, x an eigenvariable of this inference: in particular, it is not free in y, s , and Γ', Δ' . The idea in [28] is to combine the above rules in a system, with the decide on A occurring below any occurrences of the decide on $B(s, w)$, and the decide on A is turned into a “silent rule” that adds the eigenvariable w to the signature of $\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta$:

$$\frac{x \in y, \Gamma' \uparrow \cdot \vdash \cdot \uparrow x \in s, \Delta' \quad y \in w, \Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} \quad \vdots \quad \frac{\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta}{\Gamma \uparrow \cdot \vdash \cdot \uparrow \Delta} \text{ provided that } w \text{ is new.}$$

²The same kind of characterization is present in [7] in the setting of axioms as *structural* rules over *propositional* intuitionistic linear logic.

In this compound inference rule, the assumption $B(s, w)$ is not written as an assumption in the sequent but rather as a synthetic rule that is allowed only above the lower inference rule.

Another interesting example is the following quantificational instance of the axiom $\neg\alpha \vee \neg\neg\alpha$ considered in [30].

Example 22. Let B be the polarized formula $\forall x[(P(x) \supset f^+) \vee^+ ((P(x) \supset f^+) \supset f^+)]$, where the atomic formula $P(x)$ is positive. The derivation

$$\frac{\frac{\frac{\Gamma' \vdash P(t) \Downarrow \Delta'}{\Gamma' \Downarrow P(t) \supset f^+ \vdash \Delta'}}{\vdots} \quad \frac{\frac{\Gamma'', P(t) \Uparrow \cdot \vdash \cdot \Uparrow \Delta''}{\Gamma'' \Downarrow (P(t) \supset f^+) \supset f^+ \vdash \Delta''}}{\vdots} \quad \frac{\Gamma, P(t) \supset f^+ \Uparrow \cdot \vdash \cdot \Uparrow \Delta \quad \Gamma, (P(t) \supset f^+) \supset f^+ \Uparrow \cdot \vdash \cdot \Uparrow \Delta}{\Gamma \Uparrow (P(t) \supset f^+) \vee^+ ((P(t) \supset f^+) \supset f^+) \vdash \cdot \Uparrow \Delta}}{\Gamma \Downarrow \forall x(P(x) \supset f^+) \vee^+ ((P(x) \supset f^+) \supset f^+) \vdash \Delta}$$

justifies the system of rules

$$\frac{\frac{\frac{\Gamma', P(t) \vdash \Delta'}{\vdots} \quad \frac{\Gamma'', P(t) \vdash \Delta''}{\Gamma'' \vdash \Delta''}}{\Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta}{\Gamma \vdash \Delta}}{\Gamma \vdash \Delta}$$

The system of rules above corresponds exactly to the 2-system derived in [30] by translating the hypersequent rule equivalent to the (propositional version of) B .

5. Labeled proof systems for propositional modal logics

In this section, we show how to similarly apply focused proof systems LKF and LJF to the proof theory of propositional modal logics.

Following the same lines as in Section 2.4 for first-order systems, we shall show how to emulate precisely rules for modalities in labeled modal systems as synthetic connectives. Such tight emulation means that if one does focused proof search or proof checking on the polarized first-order translation of modal formulas, one is modeling nothing more or less than proof search and proof checking in the corresponding modal labeled system. As a result, we are able to show that we can use focused proofs to precisely emulate modal proofs whenever Kripke frames are characterized by bipolar properties.

This section is an extended version of [13]. While in that earlier work only (classical) modal systems from [11] were addressed, we show here that different classical and intuitionistic modal systems present in the literature can simply be computed using both polarization and focusing.

5.1. Modal logic

The language of (*propositional, normal*) modal formulas consists of a denumerable set \mathcal{P} of *propositional symbols* and a complete base of propositional connectives enhanced with the unary *modal operators* \Box and \Diamond concerning necessity and possibility, respectively.

The semantics of modal logics is often described using *Kripke models*. Here, we will follow the approach in [5], where a modal logic is defined directly by the first-order formulas capturing its intended Kripke semantics. However, here, we will map modal formulas into polarized formulas in *LKF* by generalizing the left/right translation $[\cdot]^{l/r}$ from Section 2.4. Formally, given a first-order variable x (intended to range over worlds), we define the translations $[\cdot]_x^r$ and $[\cdot]_x^l$ from modal formulas into polarized first-order formulas as follows: if P is propositional symbol, then $[P]_x^r = [P]_x^l = P(x)$ (where P is also consider a predicate symbol in first-order logic); if the top-level symbol of A is a logical connective or quantifier, then $[A]_x^{l/r}$ mimics the same translation as $[A]^{l/r}$; and if the top-level symbol is a modal operator, then we have

$$\begin{aligned} [\Box A]_x^l &= \forall y(R(x, y) \supset \partial_+([A]_y^l)) & [\Box A]_x^r &= \partial_+(\forall y(R(x, y) \supset [A]_y^r)) \\ [\Diamond A]_x^l &= \partial_-(\exists y(R(x, y) \wedge^+ [A]_y^l)) & [\Diamond A]_x^r &= \exists y(R(x, y) \wedge^+ \partial_-([A]_y^r)) \end{aligned}$$

where $R(x, y)$ is a binary predicate (that denotes the accessibility relation in a Kripke frame). Thus, following well-known characterizations of modal logic (see, for example, [5]), we know that

$$\vdash_K A \quad \text{if and only if} \quad \vdash_{LKF} \forall x.[A]_x^r$$

where K is proof system for basic classical modal logic, and we know that

$$\vdash_{IK} A \quad \text{if and only if} \quad \vdash_{LJF} \forall x.[A]_x^r$$

where IK is proof system for basic intuitionistic modal logic.

Several additional modal logics can be defined as extensions of K or IK by simply restricting the class of frames we consider. Many of the restrictions we are interested in are definable as formulas of first-order logic over the binary predicate $R(x, y)$ which encodes the accessibility relation. Table 1 contains some common restrictions on frames by listing the modal axiom capturing them together with the corresponding first-order formulas used to restrict the frame relation in the classical setting [31]. This is also true for any extension of LJF by path axioms plus contrapositives w.r.t. their corresponding models [32, 5].

5.2. Labeled proof systems for modal logics

The idea behind labeled proof systems for modal logic is to internalize elements of the corresponding Kripke semantics (namely, the worlds of a Kripke structure and the accessibility relation between such worlds) into the syntax and proof rules. As concrete examples of such a system, here we will consider the modal systems presented in [5, 12, 11]. *Labeled modal formulas* are either *labeled*

Axiom	Condition	First-Order Formula
T: $\Box A \supset A$	Reflexivity	$\forall x.R(x, x)$
4: $\Box A \supset \Box \Box A$	Transitivity	$\forall x, y, z.(R(x, y) \wedge R(y, z)) \supset R(x, z)$
5: $\Box A \supset \Box \Diamond A$	Euclideaness	$\forall x, y, z.(R(x, y) \wedge R(x, z)) \supset R(y, z)$
B: $A \supset \Box \Diamond A$	Symmetry	$\forall x, y.R(x, y) \supset R(y, x)$
3: $\Box(\Box A \supset B) \vee \Box(\Box B \supset A)$	Connectedness	$\forall x, y, z.(R(x, y) \wedge R(x, z)) \supset (R(y, z) \vee R(z, y))$
D: $\Box A \supset \Diamond A$	Seriality	$\forall x \exists y.R(x, y)$
2: $\Diamond \Box A \supset \Box \Diamond A$	Directedness	$\forall x, y, z.(R(x, y) \wedge R(x, z)) \supset \exists t(R(y, t) \wedge R(z, t))$

Table 1: Axioms and corresponding first-order conditions on R .

formulas of the form $x : A$ or *relational atoms* of the form xRy , where x and y range over a set of variables and A is a modal formula. In the following, we will use φ and ψ to denote labeled modal formulas. *Labeled sequents* have the form $\Gamma \vdash \Delta$, where Γ, Δ are multisets containing labeled modal formulas and where Δ has the usual restriction of containing at most one formula in the intuitionistic case.

In Figure 5, we present the propositional rules and some modal rules for the core labeled *classical* modal calculus. The additional modal rules for systems $G3K$ [11] and $S(K)$ [12] are depicted in Figures 6a and 6b, respectively³. The rules of the intuitionistic modal system $\mathcal{L}_{\Box\Diamond}$ [5] correspond to the rules of $G3K$ with the restriction that the consequent may have at most one *labeled* formula. In an intuitionistic version of $S(K)$, sequents have the restriction of having at most one *labeled modal* formula (there is a brief discussion of an intuitionistic version of $S(K)$ in [12], Sec. 6.2 end of page 148).

5.3. From labeled modal formulas to polarized first-order formulas

The translation $[\cdot]^{l/r}$ from labeled modal formulas into polarized first-order formulas is defined as $[x : A]^{l/r} = [A]_x^{l/r}$ and $[xRy]^{l/r} = R(x, y)$. In the following, we will sometimes use the natural extension of this notion to multisets of labeled formulas.

Finally, we define a translation from labeled sequents into focused sequents

$$[(\varphi_1, \dots, \varphi_n \vdash \psi_1, \dots, \psi_m)] = [\varphi_1]^l, \dots, [\varphi_n]^l \uparrow \cdot \vdash \cdot \uparrow [\psi_1]^r, \dots, [\psi_m]^r$$

with the restriction of $m \leq 1$ for LJF .

The results of Lemmas 3 and 5 and Theorems 4 and 6 can be then easily transported to the modal case.

³Observing that we adopt additive rules for conjunction and disjunction, multiplicative rules for implication and explicit contraction.

INITIAL RULES

$$\frac{}{x : P, \Gamma \vdash \Delta, x : P} \textit{init} \quad \frac{}{xRy, \Gamma \vdash \Delta, xRy} \textit{init}_R$$

STRUCTURAL RULES

$$\frac{\varphi, \varphi, \Gamma \vdash \Delta}{\varphi, \Gamma \vdash \Delta} C_l \quad \frac{\Gamma \vdash \Delta, \psi, \psi}{\Gamma \vdash \Delta, \psi} C_r$$

PROPOSITIONAL RULES

$$\frac{x : A, \Gamma \vdash \Delta}{x : A \wedge B, \Gamma \vdash \Delta} L\wedge_1 \quad \frac{x : B, \Gamma \vdash \Delta}{x : A \wedge B, \Gamma \vdash \Delta} L\wedge_2 \quad \frac{\Gamma \vdash \Delta, x : A \quad \Gamma \vdash \Delta, x : B}{\Gamma \vdash \Delta, x : A \wedge B} R\wedge$$

$$\frac{x : A, \Gamma \vdash \Delta \quad x : B, \Gamma \vdash \Delta}{x : A \vee B, \Gamma \vdash \Delta} L\vee \quad \frac{\Gamma \vdash \Delta, x : A}{\Gamma \vdash \Delta, x : A \vee B} R\vee_1 \quad \frac{\Gamma \vdash \Delta, x : B}{\Gamma \vdash \Delta, x : A \vee B} R\vee_2$$

$$\frac{\Gamma \vdash \Delta_1, x : A \quad x : B, \Gamma \vdash \Delta_2}{x : A \supset B, \Gamma \vdash \Delta_1, \Delta_2} L\supset \quad \frac{x : A, \Gamma \vdash \Delta, x : B}{\Gamma \vdash \Delta, x : A \supset B} R\supset$$

$$\frac{}{x : f, \Gamma \vdash \Delta} f \quad \frac{}{\Gamma \vdash \Delta, x : t} t$$

MODAL RULES

$$\frac{xRy, \Gamma \vdash \Delta, y : A}{\Gamma \vdash \Delta, x : \Box A} R\Box \quad \frac{xRy, y : A, \Gamma \vdash \Delta}{x : \Diamond A, \Gamma \vdash \Delta} L\Diamond$$

Figure 5: Some classical labeled rules, where P is an atomic formula and the eigenvariable y does not occur free in any formula of the conclusion of rules $R\Box$ and $L\Diamond$.

Theorem 23. *Let Π be a G3K derivation of a sequent S from the sequents S_1, \dots, S_n . Assume that the predicate $R(x, y)$ has positive polarity. Then there exists an LKF derivation Π' of $[S]$ from $[S_1], \dots, [S_n]$ (such that each rule application in Π corresponds to a synthetic inference rule in Π'). The exact same statement holds for $\mathcal{L}_{\Box\Diamond}$ and LJF. For $S(K)$, it holds when $R(x, y)$ has negative polarity.*

PROOF. The proof proceeds exactly as in Lemma 3 and Theorem 4, so we will show only the classical cases involving modal connectives. For the modal rules in the core fragment, the translation of the $R\Box$ from Figure 5 is given by following derivation in LKF

$$\frac{\frac{\frac{\Gamma', R(x, y) \uparrow \cdot \vdash \cdot \uparrow [A]_y^r, \Delta'}{\Gamma' \uparrow R(x, y) \vdash [A]_y^r \uparrow \Delta'} S_t, S_r}{\Gamma' \uparrow \cdot \vdash \forall y (R(x, y) \supset [A]_y^r) \uparrow \Delta'} \forall_r, \supset_r}{\Gamma' \vdash \partial_+ (\forall y (R(x, y) \supset [A]_y^r)) \downarrow \Delta'} \partial_+^r, R_r}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} D_r$$

$$\begin{array}{c}
\frac{y : A, xRy, \Gamma \vdash \Delta}{x : \Box A, xRy, \Gamma \vdash \Delta} L\Box_3 \\
\frac{xRy, \Gamma \vdash \Delta, y : A}{xRy, \Gamma \vdash \Delta, x : \Diamond A} R\Diamond_3 \\
\text{(a) Labeled modal rules for } G3K
\end{array}
\qquad
\begin{array}{c}
\frac{\Gamma \vdash \Delta_1, xRy \quad y : A, \Gamma \vdash \Delta_2}{x : \Box A, \Gamma \vdash \Delta_1, \Delta_2} L\Box_s \\
\frac{\Gamma \vdash \Delta, xRy \quad \Gamma \vdash \Delta, y : A}{\Gamma \vdash \Delta, x : \Diamond A} R\Diamond_s \\
\text{(b) Labeled modal rules for } S(K)
\end{array}$$

Figure 6: Labeled systems for the logic K .

where Γ' is any extension of $[\Gamma]$ and Δ' is any extension of $[\Delta]$ containing the formula $[x : \Box A]^r = \partial_+(\forall y(R(x, y) \supset [A]_y^r))$. Note that the condition on free variables in the definition of extension ensures that \forall can be applied in the derivation above, as the constraint on eigenvariables is satisfied. Note also that the polarity of $R(x, y)$ does not affect the shape of the derivation. The case for $L\Diamond$ is analogous.

For the distinguished modal rules in Figure 6, consider the derivation

$$\frac{\frac{\Gamma' \vdash R(x, y) \Downarrow \Delta'}{\Gamma' \Downarrow R(x, y) \supset \partial_+([A]_y^l) \vdash \Delta'} \supset_l \quad \frac{\Gamma, [A]_y^l \Uparrow \cdot \vdash \cdot \Uparrow \Delta'}{\Gamma' \Downarrow \partial_+([A]_y^l) \vdash \Delta'} R_l, \partial_+^l, S_l}{\frac{\Gamma' \Downarrow \forall y(R(x, y) \supset \partial_+([A]_y^l)) \vdash \Delta'}{\Gamma' \Uparrow \cdot \vdash \cdot \Uparrow \Delta'} \forall_l \quad D_l}$$

where Γ' is any extension of $[\Gamma]$ containing $[x : \Box A]^l = \forall y(R(x, y) \supset \partial_+([A]_y^l))$ and Δ' is any extension of $[\Delta]$. Now, if the polarity of $R(x, y)$ is positive, then π consists of the application of I_r , $R(x, y)$ should occur in Γ and the synthetic inference rule is the translation of the rule $L\Box_3$ presented in Figure 6a.

If $R(x, y)$ has, instead, negative polarity, then focus is lost in π and $R(x, y)$ is stored in the right context, producing the border sequent $\Gamma' \Uparrow \cdot \vdash \cdot \Uparrow R(x, y), \Delta'$. Hence the synthetic inference rule is the translation of the rule $L\Box_s$ presented in Figure 6b.

Finally, the initial rule

$$\frac{}{xRy, \Gamma \vdash \Delta, xRy} \text{init}_R$$

has corresponding synthetic rules in LKF

$$\frac{\Gamma' \vdash R(x, y) \Downarrow \Delta'}{\Gamma' \Uparrow \cdot \vdash \cdot \Uparrow \Delta'} I_r \quad D_r \qquad \frac{\Gamma' \Downarrow R(x, y) \vdash \Delta'}{\Gamma' \Uparrow \cdot \vdash \cdot \Uparrow \Delta'} I_l \quad D_l$$

depending whether $R(x, y)$ has positive or negative polarity, respectively, where Γ' is any extension of $[\Gamma]^l$ containing $R(x, y)$ and Δ' is any extension of $[\Delta]^r$ containing $R(x, y)$.

Theorem 24. *Let Π' be a proof of a sequent S' in LKF such that $S' = [S]$ for some G3K-sequent S . Assume that the predicate $R(x, y)$ has positive polarity. Then there exists a proof Π of S in G3K such that each synthetic inference rule in Π' corresponds to a single rule application in Π . The exact same statement holds for $\mathcal{L}_{\square\Diamond}$ and LJF. For $S(K)$, it holds when $R(x, y)$ has negative polarity.*

PROOF. Let T' be a border sequent such that each formula in T' is the translation of some G3K/S(K) formula and assume that T' is the conclusion of a synthetic inference rule Ξ in LKF with border sequent premises T'_i , $1 \leq i \leq 2$. Observe that the last rule applied in Ξ should be necessarily a decide rule. We claim that there exists G3K/S(K) sequents T, T_i , $1 \leq i \leq 2$ such that T'/T'_i is a contraction of $[T]/[T_i]$ and T_i are the premises of an inference rule in G3K/S(K) with conclusion T . The proof is done by case analysis on all possible G3K/S(K) formula φ on the translation of which a decide is applied. Suppose that one such cases is a D_r rule on $\varphi = x : \Diamond A$. Assume that $[x : \Diamond A]^r \in \Delta'$. Then we have the following synthetic inference rule in LKF

$$\frac{\frac{\frac{\Gamma' \uparrow \cdot \vdash \cdot \uparrow [A]_y^r, \Delta'}{\Gamma' \uparrow \cdot \vdash [A]_y^r \uparrow \Delta'} S_r}{\Gamma' \vdash \partial_-([A]_y^r) \downarrow \Delta'} R_r, \partial_-^r}{\frac{\Gamma' \vdash \exists y(R(x, y) \wedge^+ \partial_-([A]_y^r)) \downarrow \Delta'}{\Gamma' \uparrow \cdot \vdash \cdot \uparrow \Delta'} D_r} \exists_r, \wedge_r^+$$

If $R(x, y)$ is positive, then π consists of the rule l_r and Γ' must contain the formula $R(x, y)$, corresponding to the rule application of $R\Diamond_3$ in Figure 6a. If $R(x, y)$ has negative polarity, then focus is lost in π and $R(x, y)$ is stored in the right context, producing the border sequent $\Gamma' \uparrow \cdot \vdash \cdot \uparrow R(x, y), \Delta'$, corresponding to an application of the rule $R\Diamond_S$ presented in Figure 6b. Observe that decide rules in LKF carry an implicit contraction, which is discarded in G3K/S(K) by the contractions on translations.

We can now prove the theorem by proceeding top-down, starting from the leaves of Π' and building Π by repeatedly applying the method above described. At each step, we get as the conclusion of a G3K/S(K) rule application a sequent S^* such that the one obtained in the corresponding step of Π' is a contraction of $[S^*]$.

The strong correspondence between labeled rule applications and LKF/LJF synthetic inference rules can also be used to get an immediate proof of the completeness of G3K, S(K) and $\mathcal{L}_{\square\Diamond}$.

Corollary 25. *The systems G3K and S(K) (and $\mathcal{L}_{\square\Diamond}$ respectively) are complete with respect to modal logic K (IK respectively).*

PROOF. Follows from the completeness of LKF/LJF and Theorem 24.

5.4. Labeled systems for extensions of K and IK

While [12] presents a detailed study of extensions of the modal system $S(K)$ with Horn relational theories (see Section 4.2), in [5, 11] the systems $G3K$ and $\mathcal{L}_{\square\lozenge}$ were extended in order to capture theories given by geometric frame conditions. In all such works, the idea was to modularly add, to a base modal system, rules defined from the axioms according to a proper chosen scheme.

In the view of Theorem 12, different polarizations⁴ of the frame conditions presented in Table 1 correspond to different synthetic inference rules, and the statements of Theorems 23 and 24 hold also for the corresponding bipolar extensions of $G3K/S(K)/\mathcal{L}_{\square\lozenge}$.

In Figure 7 we present the relational rules capturing frame properties of Table 1, where \wedge, \vee are polarized using the positive polarity. The rules in the first column are derived when R is given a positive polarity and correspond to the geometric rule scheme presented in [5, 11]. For the rules in the second column, R was given negative polarity and they correspond to the un-skolemized version of the rules appearing in [12].

Observe that the unfocused rule corresponding to the possible bipolars for Axiom D does not depend on the polarization of R , having the form

$$\frac{R(x, y), \Gamma \vdash \Delta}{\Gamma \vdash \Delta} D$$

where $y \notin \Gamma, \Delta$.

Axiom 3 is a geometric implication but not a Horn clause, hence it is not considered in [12]. Axiom 2 is also not a Horn clause, but it can be transformed, using skolemization (see Example 20), into a set of Horn formulas, and it is considered in [12] under the name *convergency*.

The geometric and right universal rule schemes for 2 and 3 are given by

$$\frac{\frac{R(y, u), R(z, u), \Gamma \vdash \Delta}{R(x, y), R(x, z), \Gamma \vdash \Delta} 2_{GRS}}{\Gamma \vdash \Delta, R(x, y) \quad \Gamma \vdash \Delta, R(x, z) \quad R(y, u), R(z, u), \Gamma \vdash \Delta}{\Gamma \vdash \Delta} 2_{RR}$$

where u does not occur in the conclusion.

$$\frac{\frac{R(y, z), \Gamma \vdash \Delta \quad R(z, y), \Gamma \vdash \Delta}{R(x, y), R(x, z), \Gamma \vdash \Delta} 3_{GRS}}{\frac{R(y, z), \Gamma \vdash \Delta \quad R(z, y), \Gamma \vdash \Delta \quad \Gamma \vdash \Delta, R(x, y) \quad \Gamma \vdash \Delta, R(x, z)}{\Gamma \vdash \Delta} 3_{RR}}$$

⁴Remembering that, in the directedness axiom, \wedge in the head of the clause should necessarily be translated to \wedge^+ . Any polarization of the other axioms results on bipolar formulas.

Axiom	Geometric rule scheme	Right universal rule scheme
T	$\frac{R(x, x), \Gamma \vdash \Delta}{\Gamma \vdash \Delta} T_{GRS}$	$\frac{}{\Gamma \vdash \Delta, R(x, x)} T_{RR}$
4	$\frac{R(x, z), \Gamma \vdash \Delta}{R(x, y), R(y, z), \Gamma \vdash \Delta} 4_{GRS}$	$\frac{\Gamma \vdash \Delta, R(x, y) \quad \Gamma \vdash \Delta, R(y, z)}{\Gamma \vdash \Delta, R(x, z)} 4_{RR}$
5	$\frac{R(y, z), \Gamma \vdash \Delta}{R(x, y), R(x, z), \Gamma \vdash \Delta} 5_{GRS}$	$\frac{\Gamma \vdash \Delta, R(x, y) \quad \Gamma \vdash \Delta, R(x, z)}{\Gamma \vdash \Delta, R(y, z)} 5_{RR}$
B	$\frac{R(y, x), \Gamma \vdash \Delta}{R(x, y), \Gamma \vdash \Delta} B_{GRS}$	$\frac{\Gamma \vdash \Delta, R(x, y)}{\Gamma \vdash \Delta, R(y, x)} B_{RR}$

Figure 7: Axioms and corresponding sequent rules.

6. Conclusion

We have described how the notion of synthetic inference rule that is provided by sequent calculus notions of polarization and focusing can be used to provide inference rules that capture certain classes of axioms. In particular, focused proof systems naturally lead to the notion of bipolar formulas and these result in synthetic inference rules that only need to mention atomic formulas. We show that geometric formulas are examples of such bipolar formulas and that polarized versions of such formulas yield known inference systems derived from geometric formulas. Certain subsets of geometric formulas admit more than one polarization and these variations explain the forward-chaining and backward-chaining variants of their synthetic inference rules. The cut-elimination theorem for focused proof systems also provides a direct proof of cut-elimination for the proof systems that arise from incorporating synthetic inference rules based on polarized formulas. Additionally, all of these results work equally well in both classical and intuitionistic logics using the corresponding *LKF* and *LJF* focused proof systems. Finally, we show how to account for and generalize labeled proof systems for propositional modal logics.

References

References

- [1] V. Danos, J.-B. Joinet, H. Schellinx, LKT and LKQ: sequent calculi for second order logic based upon dual linear decompositions of classical implication, in: J.-Y. Girard, Y. Lafont, L. Regnier (Eds.), *Advances in Linear Logic*, no. 222 in London Mathematical Society Lecture Note Series, Cambridge University Press, 1995, pp. 211–224. [3](#), [6](#)
- [2] C. Liang, D. Miller, [Focusing and polarization in intuitionistic logic](#), in: J. Duparc, T. A. Henzinger (Eds.), *CSL 2007: Computer Science Logic*, Vol.

4646 of LNCS, Springer, 2007, pp. 451–465.

URL <http://www.lix.polytechnique.fr/Labo/Dale.Miller/papers/cs107liang.pdf> 3, 6

- [3] C. Liang, D. Miller, [Focusing and polarization in linear, intuitionistic, and classical logics](#), *Theor. Comput. Sci.* 410 (46) (2009) 4747–4768. doi: [10.1016/j.tcs.2009.07.041](https://doi.org/10.1016/j.tcs.2009.07.041).
URL <http://dx.doi.org/10.1016/j.tcs.2009.07.041> 3, 6, 10, 12, 13, 20
- [4] K. Chaudhuri, [Focusing strategies in the sequent calculus of synthetic connectives](#), in: I. Cervesato, H. Veith, A. Voronkov (Eds.), *Logic for Programming, Artificial Intelligence, and Reasoning*, 15th International Conference, LPAR 2008, Doha, Qatar, November 22–27, 2008. Proceedings, Vol. 5330 of *Lecture Notes in Computer Science*, Springer, 2008, pp. 467–481. doi: [10.1007/978-3-540-89439-1_33](https://doi.org/10.1007/978-3-540-89439-1_33).
URL https://doi.org/10.1007/978-3-540-89439-1_33 3
- [5] A. K. Simpson, *The proof theory and semantics of intuitionistic modal logic*, Ph.D. thesis, College of Science and Engineering, School of Informatics, University of Edinburgh (1994). 4, 21, 28, 29, 33
- [6] R. Dyckhoff, S. Negri, [Geometrisation of first-order logic](#), *The Bulletin of Symbolic Logic* 21 (2) (2015) 123–163.
URL <http://www.jstor.org/stable/24327109> 4, 22, 25
- [7] A. Ciabattoni, N. Galatos, K. Terui, [From axioms to analytic rules in nonclassical logics](#), in: *Proceedings of the Twenty-Third Annual IEEE Symposium on Logic in Computer Science, LICS 2008*, 24–27 June 2008, Pittsburgh, PA, USA, 2008, pp. 229–240. doi: [10.1109/LICS.2008.39](https://doi.org/10.1109/LICS.2008.39).
URL <https://doi.org/10.1109/LICS.2008.39> 4, 16, 26
- [8] A. Ciabattoni, P. Maffezioli, L. Spendier, [Hypersequent and labelled calculi for intermediate logics](#), in: D. Galmiche, D. Larchey-Wendling (Eds.), *Automated Reasoning with Analytic Tableaux and Related Methods - 22th International Conference, TABLEAUX 2013*, Nancy, France, September 16–19, 2013. Proceedings, Vol. 8123 of *Lecture Notes in Computer Science*, Springer, 2013, pp. 81–96. doi: [10.1007/978-3-642-40537-2_9](https://doi.org/10.1007/978-3-642-40537-2_9).
URL https://doi.org/10.1007/978-3-642-40537-2_9 4
- [9] S. Negri, J. von Plato, [Cut elimination in the presence of axioms](#), *Bulletin of Symbolic Logic* 4 (4) (1998) 418–435.
URL <http://www.math.ucla.edu/~asl/bsl/0404/0404-003.ps> 4, 21, 23
- [10] S. Negri, [Contraction-free sequent calculi for geometric theories with an application to Barr’s theorem](#), *Arch. Math. Log.* 42 (4) (2003) 389–401. doi: [10.1007/s001530100124](https://doi.org/10.1007/s001530100124).
URL <http://dx.doi.org/10.1007/s001530100124> 4, 21, 23

- [11] S. Negri, [Proof analysis in modal logic](#), *J. Philosophical Logic* 34 (5-6) (2005) 507–544. doi:10.1007/s10992-005-2267-3.
URL <http://dx.doi.org/10.1007/s10992-005-2267-3> 4, 21, 27, 28, 29, 33
- [12] L. Viganò, *Labelled Non-Classical Logics*, Kluwer Academic Publishers, 2000. 4, 21, 25, 28, 29, 33
- [13] D. Miller, M. Volpe, [Focused labeled proof systems for modal logic](#), in: *Logic for Programming, Artificial Intelligence, and Reasoning - 20th International Conference, LPAR-20 2015, Suva, Fiji, November 24-28, 2015, Proceedings, 2015*, pp. 266–280. doi:10.1007/978-3-662-48899-7_19.
URL http://dx.doi.org/10.1007/978-3-662-48899-7_19 4, 27
- [14] S. Marin, D. Miller, M. Volpe, [A focused framework for emulating modal proof systems](#), in: L. D. Beklemishev, S. Demri, A. Maté (Eds.), *Advances in Modal Logic 11, proceedings of the 11th conference on "Advances in Modal Logic," held in Budapest, Hungary, August 30 - September 2, 2016*, College Publications, 2016, pp. 469–488.
URL <http://www.aiml.net/volumes/volume11/Marin-Miller-Volpe.pdf> 4
- [15] G. Gentzen, *Investigations into logical deduction*, in: M. E. Szabo (Ed.), *The Collected Papers of Gerhard Gentzen*, North-Holland, Amsterdam, 1935, pp. 68–131. doi:10.1007/BF01201353. 5
- [16] D. Miller, G. Nadathur, F. Pfenning, A. Scedrov, *Uniform proofs as a foundation for logic programming*, *Annals of Pure and Applied Logic* 51 (1–2) (1991) 125–157. 6
- [17] D. Miller, G. Nadathur, A. Scedrov, *Hereditary Harrop formulas and uniform proof systems*, in: D. Gries (Ed.), *2nd Symp. on Logic in Computer Science*, Ithaca, NY, 1987, pp. 98–105. 6
- [18] J. Andreoli, [Logic programming with focusing proofs in linear logic](#), *J. Log. Comput.* 2 (3) (1992) 297–347. doi:10.1093/logcom/2.3.297.
URL <http://dx.doi.org/10.1093/logcom/2.3.297> 6
- [19] H. Herbelin, *A lambda-calculus structure isomorphic to Gentzen-style sequent calculus structure*, in: *Computer Science Logic, 8th International Workshop, CSL '94, Vol. 933 of Lecture Notes in Computer Science*, Springer, 1995, pp. 61–75. 6
- [20] R. Dyckhoff, S. Lengrand, *LJQ: a strongly focused calculus for intuitionistic logic*, in: A. Beckmann, *et al.* (Eds.), *Computability in Europe 2006*, Vol. 3988 of LNCS, Springer, 2006, pp. 173–185. 6
- [21] C. Liang, D. Miller, *Focusing Gentzen's LK proof system*, submitted (Apr. 2020). 10, 20, 21

- [22] V. Nigam, D. Miller, [A framework for proof systems](#), *J. Autom. Reasoning* 45 (2) (2010) 157–188. doi:10.1007/s10817-010-9182-1. URL <https://doi.org/10.1007/s10817-010-9182-1> 15
- [23] D. Miller, G. Nadathur, *Programming with Higher-Order Logic*, Cambridge University Press, 2012. doi:10.1017/CB09781139021326. 19, 38
- [24] G. Nadathur, D. Miller, [An Overview of \$\lambda\$ Prolog](#), in: *Fifth International Logic Programming Conference*, MIT Press, Seattle, 1988, pp. 810–827. URL <http://www.lix.polytechnique.fr/Labo/Dale.Miller/papers/iclp88.pdf> 19, 38
- [25] J.-Y. Girard, *Proof Theory and Logical Complexity*, Vol. I of *Studies in Proof Theory*, Bibliopolis, edizioni di filosofia e scienze, Napoli, 1987. 21
- [26] R. Ramanayake, [From axioms to proof rules, then add quantifiers](#), in: C. Benzmüller, J. Otten (Eds.), *Proceedings of the 2nd International Workshop Automated Reasoning in Quantified Non-Classical Logics (ARQNL 2016) affiliated with the International Joint Conference on Automated Reasoning (IJCAR 2016)*, Coimbra, Portugal, July 1, 2016, Vol. 1770 of *CEUR Workshop Proceedings*, CEUR-WS.org, 2016, pp. 1–8. URL http://ceur-ws.org/Vol-1770/ARQNL2016_invited1.pdf 21
- [27] S. Negri, J. von Plato, [Proof Analysis - a contribution to Hilbert’s last problem](#), Cambridge University Press, 2011. doi:10.1017/CB09781139003513. URL <http://www.cambridge.org/de/academic/subjects/philosophy/logic/proof-analysis-contribution-hilberts-last-problem?format=PB> 21, 23
- [28] S. Negri, [Proof analysis beyond geometric theories: from rule systems to systems of rules](#), *J. Log. Comput.* 26 (2) (2016) 513–537. doi:10.1093/logcom/exu037. URL <http://dx.doi.org/10.1093/logcom/exu037> 23, 24, 26
- [29] E. J. Lemmon, D. S. Scott, *An Introduction to Modal Logic*, Blackwell, 1977. 25
- [30] A. Ciabattoni, F. A. Genco, [Hypersequents and systems of rules: Embeddings and applications](#), *ACM Trans. Comput. Log.* 19 (2) (2018) 11:1–11:27. doi:10.1145/3180075. URL <https://doi.org/10.1145/3180075> 26, 27
- [31] H. Sahlqvist, [Completeness and correspondence in first and second order semantics for modal logic](#), in: N. H. S. Kanger (Ed.), *Proceedings of the Third Scandinavian Logic Symposium*, 1975, pp. 110–143. 28
- [32] G. D. Plotkin, C. P. Stirling, [A framework for intuitionistic modal logic](#), in: J. Y. Halpern (Ed.), *1st Conference on Theoretical Aspects of Reasoning About Knowledge*, Morgan Kaufmann, 1986, pp. 399–406. 28

Appendix A. Computing bipoles from bipolar formulas

We briefly describe a logic program that can compute bipole inference rules from bipolar formulas: this implementation uses the λ Prolog programming language [23, 24]. The key novelties of the syntax of λ Prolog for this implementation are the following. Since this logic programming language is typed, the keyword `kind` is used to introduce a new primitive type, and the keyword `type` is used to introduce new typed constructors. Also, the backslash `\` is an infix binding construction: for example, the expression `pi x \ B` denotes the universal quantification of `x` over the formulas `B`. This same backslash operator is also used to build λ -bindings within terms.

The datatype of polarized formulas is given by the following declarations.

```
kind i, atm, fm                                type.
type atm                                       atm -> fm.
type all, some                                 (i -> fm) -> fm.
type imp, pand, nand, por, nor                fm -> fm -> fm.
type ptrue, ntrue, pfalse, nfalse            fm.
```

Formulas and atoms are given a polarity using the following declarations and clauses. Here, the predicate `delta` encodes an atomic bias assignment (see Section 2.2).

```
kind bias                                       type.
type pp, nn                                    bias.
type delta                                     atm -> bias -> o.

type neg, pos                                  fm -> o.
type patom, natom                             fm -> o.
```

```
patom (atm A) :- delta A pp.
natom (atm A) :- delta A nn.
```

```
neg (imp _ _) & neg (all _).
neg (nand _ _) & neg (nor _ _) & neg ntrue & neg nfalse.
neg A :- natom A.
pos (some _).
pos (pand _ _) & pos (por _ _) & pos ptrue & pos pfalse.
pos A :- patom A.
```

The various sequents are encoded using the following declarations and clauses.

```
kind premise                                   type.
type truep                                     premise.
type andp                                       premise -> premise -> premise.
type allp                                       (i -> premise) -> premise.
type primitivep                               premise -> o.
type borders                                  premise -> o.

kind rhs                                       type.
type rl, rr                                    fm -> rhs.
```

```

type async          list fm -> list fm -> rhs -> premise.
type syncR         list fm -> fm -> premise.
type syncL         list fm -> fm -> fm -> premise.

primitivep (async _ _ _).
primitivep (syncL _ _ _) & primitivep (syncR _ _).

borders (async _ [] (rr _)).
borders truep.
borders (andp P1 P2) :- borders P1, borders P2.
borders (allp P) :- pi x\ borders (P x).

```

The following four sequent structures are encoded using the corresponding λ Prolog terms, all four of which are also considered to be primitive premises (using the specification for `primitivep` above).

$$\begin{array}{l|l}
\Gamma \uparrow \Theta \vdash E \uparrow \cdot & (\text{async } \Gamma \Theta \text{ (rl } E)) \\
\Gamma \uparrow \Theta \vdash \cdot \uparrow E & (\text{async } \Gamma \Theta \text{ (rr } E)) \\
\Gamma \downarrow B \vdash E & (\text{syncL } \Gamma B E) \\
\Gamma \vdash B \downarrow E & (\text{syncR } \Gamma B E)
\end{array}$$

Figure A.8 encodes the various rules of *LJF* (except for the decide rules) using the binary predicate `rule`. Figure A.9 defines three predicates: `rotate` ensures that tree-structure of primitive premises is organized more as a list; `red1` holds if exactly one inference rule is applied to exactly one premise in its first argument; and `reduce` repeatedly applies `red1` until only border sequents remain. Consider proving the goal

```
reduce (syncL Gamma F (atm B)) Premises.
```

for different instantiations of the variable `F` and for different polarity assumptions on atomic formulas. First, assume that all atomic formulas are positive. If `F` is instantiated with the term

```
(all u\ all v\ all w\ imp (atm (adj u v))
                        (imp (atm (path v w)) (atm (path u w))))),
```

which encodes the formulas $\forall u \forall v \forall w (adj\ u\ v \supset path\ v\ w \supset path\ u\ w)$ then λ Prolog will solve this goal formula by computing the following substitution.

```
Gamma      = atm (adj X Z) :: atm (path Z Y) :: L
Premises   = async (atm (path X Y) :: atm (adj X Z) ::
                  atm (path Z Y) :: L) nil (rr (atm B))
```

The inference rule computed by solving this query is

$$\frac{adj\ X\ Z, path\ Z\ Y, path\ X\ Y, L \uparrow \cdot \vdash \cdot \uparrow B}{adj\ X\ Z, path\ Z\ Y, L \uparrow \cdot \vdash \cdot \uparrow B}$$

Here, `X`, `Y`, and `Z` are schema variables of type `i`, `L` is a schema variable of type `list fm`, and `B` is a schema variable of type `fm`. Next, assume that all atomic formulas are negative. Executing the same goal as before (for the same instantiation for `F`) yields the substitution

```

type rule          premise -> premise -> o.
rule (async Gm [ptrue|Th] R) (async Gm Th R).
rule (async Gm [pand B C|Th] R) (async Gm [B,C|Th] R).
rule (async Gm [pfalse|Th] R) truep.
rule (async Gm [por B C|Th] R) (andp (async Gm [B|Th] R)
                                   (async Gm [C|Th] R)).
rule (async Gm [some B|Th] R) (allp x\ async Gm [B x|Th] R).
rule (async Gm [C|Th] R) (async (C::Gm) Th R) :-
                                   neg C; patom C.
rule (async Gm [] (rl (nand B C)))(andp (async Gm [] (rl B))
                                         (async Gm [] (rl C))).
rule (async Gm [] (rl ntrue)) truep.
rule (async Gm [] (rl (imp B C))) (async Gm [B] (rl C)).
rule (async Gm [] (rl (all B)))
    (allp x\ async Gm [] (rl (B x))).
rule (async Gm [] (rl D)) (async Gm [] (rr D)) :-
                                   pos D; natom D.

rule (syncL Gm (all B) R) (syncL Gm (B T) R).
rule (syncL Gm A A) truep :- natom A.
rule (syncL Gm (imp B C) R)
    (andp (syncR Gm B) (syncL Gm C R)).
rule (syncL Gm (nand B C) R) (syncL Gm B R).
rule (syncL Gm (nand B C) R) (syncL Gm C R).
rule (syncL Gm P R) (async Gm [P] (rr R)) :- pos P.
rule (syncR Gm A) truep :- patom A, memb A Gm.
rule (syncR Gm ptrue) truep.
rule (syncR Gm (pand B C)) (andp (syncR Gm B) (syncR Gm C)).
rule (syncR Gm (por B C)) (syncR Gm B).
rule (syncR Gm (por B C)) (syncR Gm C).
rule (syncR Gm (some B)) (syncR Gm (B T)).
rule (syncR Gm N) (async Gm [] (rl N)) :- neg N.

```

Figure A.8: The predicate rule encodes the rules of *LJF* (Figure 3) except for D_l and D_r .

```

type rotate, red1, reduce    premise -> premise -> o.
rotate Prim Prim :- primitivep Prim.
rotate truep truep.
rotate (andp R S) (andp R T) :- primitivep R, rotate S T.
rotate (andp truep U) T :- rotate U T.
rotate (andp (andp R S) U) T :- rotate (andp R (andp S U)) T.
rotate (andp (allp R) S) T :- rotate (allp x\ andp (R x) S) T.
rotate (allp R) (allp T) :- pi x\ rotate (R x) (T x).
red1 G H :- primitivep G, rule G H.
red1 (andp G1 G2) (andp H G2) :- red1 G1 H, !.
red1 (andp G1 G2) (andp G1 H) :- red1 G2 H.
red1 (allp G) (allp K) :- pi x\ red1 (G x) (K x).
reduce Gs Hs :- red1 Gs Ks, rotate Ks Rs, reduce Rs Hs.
reduce Gs Gs :- borders Gs.

```

Figure A.9: Three predicated useful for computing bipoles.


```

Premises = andp (async Gamma nil (rr (atm (adj X Y))))
           (andp (async Gamma nil (rr (atm (path Y Z)))) truep)
B         = path X Z

```

Thus, the inference rule computed by solving this query is

$$\frac{\Gamma \uparrow \cdot \vdash \cdot \uparrow \text{adj } X \ Y \quad \Gamma \uparrow \cdot \vdash \cdot \uparrow \text{path } Y \ Z}{\Gamma \uparrow \cdot \vdash \cdot \uparrow \text{path } X \ Z}$$

For a final example, consider solving this goal but where **F** is instantiated to the term

```

(all u\ all v\ imp (all w\ imp (atm (in w u)) (atm (in w v)))
 (atm (subset u v))).

```

and where atomic formulas have negative bias. The computed substitution is then

```

Premises = (allp w\ andp (async (atm (in w X) :: Gamma) nil
                               (rr (atm (in w Y)))) truep)
B         = subset X Y

```

Thus, the inference rule computed by solving this query is

$$\frac{\Gamma, \text{in } w \ X \uparrow \cdot \vdash \cdot \uparrow \text{in } w \ Y}{\Gamma \uparrow \cdot \vdash \cdot \uparrow \text{subset } X \ Y}$$

Here, w is an eigenvariable for this inference figure and it corresponds to the binding `allp w\` in the answer substitution for **Premises** above.